

# 3

Human IgG3 with extended half-life does not improve Fc-gamma Receptor-mediated cancer antibody therapies in mice

**Rens Braster**<sup>1</sup>, Simran Grewal<sup>1,2</sup>, Remco Visser<sup>3</sup>, Helga K. Einarsdottir<sup>3</sup>, Marjolein van Egmond<sup>1,2</sup>, Gestur Vidarsson<sup>3,5</sup>, Marijn Bögels<sup>1,2,5</sup>

*Manuscript submitted*

## **ABSTRACT**

Current anti-cancer therapeutic antibodies that are used in the clinic are either humanized or fully human immunoglobulin G1 (IgG1). These antibodies bind with high affinity to the target antigen and are efficient in activating the immune system via IgG Fc receptors and/or complement. In addition to IgG1, three more isotypes are present in humans, of which IgG3 has been found to be superior compared to human IgG1 in inducing antibody dependent cell cytotoxicity (ADCC), phagocytosis or activation of complement in some models. Nonetheless, no therapeutic human IgG3 mAbs have been developed due to the short *in vivo* half-life of most known IgG3 allotypes. In this manuscript, we compared the efficacy of V-gene matched IgG1 and IgG3 anti-tumour mAb (TA99) in mice, using natural variants of human IgG3 with short- or long half-life, differing only at position 435 with and arginine or histidine, respectively.

*In vitro* human IgG1 and IgG3 did not show any differences in opsonisation ability of B16F10-gp75 mouse melanoma cells. IgG1, however, was superior in inducing phagocytosis of tumour cells by mouse macrophages. Similarly, in a mouse peritoneal metastasis model we did not detect an improved effect of IgG3 in preventing tumour outgrowth. Moreover, replacing the arginine at position 435 for a histidine in IgG3 to enhance half-life did not result in better suppression of tumour outgrowth compared to wild type IgG3 when injected prior to tumour cell injection.

In conclusion, human IgG3 does not have improved therapeutic efficacy compared to human IgG1 in a mouse tumour model.

<sup>1</sup>Department of Molecular Cell Biology and Immunology, VU University medical centre, Amsterdam, the Netherlands;

<sup>2</sup>Department of Surgery, VU University medical centre Amsterdam, the Netherlands;

<sup>3</sup>Department of Experimental Immunohematology, Sanquin Research and Landsteiner Laboratory, Academic Medical Centre, University of Amsterdam, Amsterdam, the Netherlands;

<sup>§</sup>Authors contributed equally to this manuscript.

## INTRODUCTION

The development of new and better monoclonal antibodies (mAb) to use as therapy to treat cancer - in addition to chemo- and/ or radiotherapy - has increased dramatically in the last decade<sup>1</sup>. mAbs can be designed to specifically interact with tumour-associated antigens, and initiate a wide range of effector mechanisms, which can potentially result in regression of the tumour. Several anti-tumour mAbs have now been approved for cancer therapy by the American food and drug administration (FDA), and the number of potential new targets is increasing rapidly. The most successful mAb that is currently used in the clinic is rituximab, which is widely employed in the treatment of several B-cell malignancies and has significantly improved clinical outcome of patients<sup>2</sup>. Additionally, anti-HER-2/neu and anti-epidermal growth factor receptor mAbs are increasingly used to treat several malignancies such as mammary carcinoma, colorectal cancer or head and neck cancer respectively. However, despite some clinical successes, a substantial proportion of cancer patients fail to achieve complete remission or experience relapse after receiving mAb therapy. Improvement of antibody immunotherapeutic approaches is therefore warranted.

Therapeutic mAbs can trigger a multitude of functions to eliminate tumour cells. These include direct effects, such as inducing growth arrest of tumour cells by blocking the binding of growth factors to their receptor, or initiation of apoptosis<sup>1,3</sup>. In addition, after binding to tumour cells they can activate the complement system to destroy the target, which is referred to as complement dependent cytotoxicity (CDC). Through their Fc part, mAbs of the immunoglobulin G (IgG) isotype can furthermore bind to IgG Fc receptors (Fc receptors) that are expressed on immune effector cells. This can lead to killing of tumour cells via a process referred to as antibody dependent cellular cytotoxicity (ADCC) or through antibody dependent cellular phagocytosis (ADCP).

In vivo, Fc $\gamma$  receptor mediated mechanisms were shown to play a prominent role in antibody-mediated (TA99) therapies against the syngeneic B16 melanoma model<sup>4</sup>. In this model, anti-tumour mAb treatment of Fc receptor  $\gamma$  chain-deficient mice, which lack the activating receptors Fc $\gamma$ RI, Fc $\gamma$ RIII and Fc $\gamma$ RIV, did not protect against the development of lung metastases, whereas control wild type mice had less tumour development after mAb therapy<sup>4</sup>. We previously demonstrated that treatment of mice with murine IgG2a (which shares functional homology with human IgG1) successfully prevented the development of liver metastases. Clinical efficacy depended on the presence of Fc $\gamma$ RI and Fc $\gamma$ RIV<sup>5</sup>. Additionally, liver macrophages (Kupffer cells) were shown to be indispensable for mAb-induced eradication of tumour cells, which was due to effective ADCP<sup>6,7</sup>.

Currently, most clinical mAb are of the IgG1 isotype, due to favourable characteristics such as the ability to induce CDC, ADCC and ADCP, combined with its long half-life. Interestingly, human IgG3 has the highest affinity for Fc $\gamma$  receptors and activates complement even more potently than IgG1<sup>8,9</sup>. Hence, IgG3 has been found to mediate even stronger effector functions than IgG1 in vitro and in mice<sup>10-12</sup>. As such, it is in theory the best ligand for all Fc $\gamma$  receptors expressed on macrophages and other immune cells<sup>13</sup>. IgG3 has however a short half-life in vivo, making it potentially less effective for mAb therapies<sup>14</sup>. It was demonstrated that human IgG1 antibodies, which are the most abundant IgG isotype in humans, inhibit FcRn-mediated transport and rescue of human IgG3, subsequently leading to lysosomal degradation of IgG3<sup>10,15</sup>. Due to the shorter half-life and earlier assumptions that the longer hinge region of IgG3 would lead to enhanced proteolytic degradation no IgG3 therapeutic antibodies are under development so far<sup>16,17</sup>. However, as it was demonstrated that



IgG3 induces superior phagocytosis of e.g. opsonised (infected) red blood cells or micro-organisms by mononuclear cells, IgG3 may represent an ideal candidate for antibody therapy to prevent the development of liver metastases<sup>18,19</sup>. We therefore developed recombinant V-gene-matched human IgG1 and IgG3 mAbs against murine B16 tumour cells, and evaluated their potential in prevention of metastases development. Additionally, we introduced an amino acid change in the Fc domain of IgG3 at position 435 that represents a natural variation also found in humans, which is essential for FcRn interactions<sup>10,20</sup>. All subclasses and IgG-species from other mammals have been found to have an histidine (H) at this position which endows IgG antibodies a pH-dependent binding to FcRn, which is crucial for their FcRn-dependent recycling properties<sup>10,21</sup>. IgG3 has classically be defined as a subclass with an arginine (R) at this position, which we recently found to be the cause of the short have-life of IgG3 due to competition with the other subclasses for FcRn-mediated transport<sup>10</sup>. However, due to an allotypic variation at this position in IgG3, some individuals actually do have histidine at this position. Unlike R435-expressing individuals, IgG3 in H435-expressing individuals has a half-life comparable to IgG1. In addition, placental transport, which is also mediated by FcRn, of IgG3 in humans is also comparable to IgG1 in H435-IgG3 expressing individuals, strongly supporting that this amino acid variation is underlying factor for the reported short half-life for IgG3<sup>22</sup>. As we and others have found IgG3 to have strong effector functions through both Fcγ receptor<sup>9,10</sup> and complement<sup>9,11,23</sup>, we now tested if H435-IgG3 has superior effector functions as therapeutic antibody (TA99) in the B16 melanoma model.

## **MATERIALS AND METHODS**

### Animals

Male C57Bl/6 mice, weighing 20-25 g, were obtained from Harlan, Boxmeer, The Netherlands. Animals were kept under standard laboratory conditions and had access to food and water ad libitum. The Committee for Animal Research of the VUmc approved all experiments, according to the European and national guidelines.

### Cell Culture

*Tumour cells.* Truncated GP75 (Tyrp-1, tyrosinase related protein 1) was transfected in mouse melanoma B16F10 (B16F10-gp75) cells as described in<sup>7</sup> and cultured in RPMI-1640 (Invitrogen, Paisley, UK), supplemented with 10% heat inactivated foetal calf serum (FCS), penicillin (100U/ml), streptomycin (100 µg/ml) and L-glutamine (200 µM) (further referred to as complete RPMI). Cell suspensions were prepared by enzymatic detachment using trypsin-EDTA solution (Invitrogen). Viability was assessed by trypan blue exclusion and always exceeded 95%.

*L929 cell conditioned medium (LCM).* Mouse L929 cells secrete macrophage- colony stimulating factor (M-CSF) and were used to produce L929 cell conditioned medium (LCM) to differentiate macrophages. L929 cells were grown to confluency, after which medium was changed with fresh complete RPMI. Cells were grown for 7 days, after which LCM was harvested, centrifuged at 4750xg for 10 minutes, filtered through 0.2 µm filters and stored at -20°C till further use.

*Murine bone marrow macrophages.* Bone marrow was harvested from freshly isolated femur, tibia and humerus from wild type C57BL/6. After removal of connective tissues and muscles, bone marrow was flushed and single cell suspensions were made by passing bone marrow through a sterile 70 µm filter (BD Falcon, Bedford, MA). Macrophages were differentiated by incubating bone marrow cells for 7 days with complete DMEM, supplemented with 15% LCM (hereafter referred to as macrophage

medium). Macrophages were harvested after 15 minute incubation with trypsin-EDTA and subsequent scraping using a cell scraper. Macrophages were seeded in 24 well plates ( $4 \times 10^5$ / well) for *in vitro* cytotoxicity assays.

#### Generation of human IgG1 and IgG3 anti-GP75 (TA99)

The variable regions of the heavy and light chains (VL, VH) were cloned from a hybridoma, which produces murine IgG2a mAb against the murine gp75 antigen. Variable regions were expressed as chimeric human IgG in a similar manner as previously described<sup>9</sup>. RNA was isolated with the RNeasy Mini Kit, (Qiagen, CA), and VH and VL genes were amplified as described in<sup>24</sup>, with CH and CL specific primers. The product was then ligated into pGEM-T (Invitrogen, CA) and sequenced by ABI 373 Stretch automated sequencing (Applied Biosystems, Foster City, CA). Codon optimized V-genes, including 5'HindIII, and 3'EcoRI restriction sites, Kozak sequence and HAVT20-leader<sup>25</sup> were then designed and ordered from MR Gene (now Geneart/Invitrogen), along with codon optimized human  $\kappa$ ,  $\gamma$ 1,  $\gamma$ 3 constant regions for the variable light and heavy chains, respectively. H/R mutations were introduced at position 435 as described by Stapleton et al<sup>10</sup>. HindIII/EcoRI fragments for the codon optimized light chain was ligated into pEE14.4 (Lonza), the HindIII/EcoRI heavy chain into pEE6.4 (Lonza). The correct assembly of the final construct was verified by sequencing, and it produced in the FreeStyle 293 expression system (Invitrogen) according to the manufacturer's instructions. Antibodies were purified on a protein A (H435 IgG) or protein G (R435 IgG) HiTrap HP column (GE Life Sciences) and dialyzed against PBS overnight.

#### Flow cytometry

Mouse B16F10 or B16F10-gp75 cell lines were incubated with different concentrations of primary mouse or human anti-gp75 mAb or human IgG1 anti-HEPC (isotype control) or mouse IgG2a anti-rat MG4 (isotype control) antibodies for 45 minutes on ice. After washing, primary antibody was detected by incubation with PE-conjugated goat-anti-human IgG mAb (1:50) or rabbit-anti-mouse IgG mAb. Cells were analysed with flow cytometry (FacsCalibur, BD, San Jose, CA).

#### Fluorescent labelling

For *in vitro* cytotoxicity assays mouse tumour cells were harvested and incubated ( $1-10 \times 10^6$  cells/ml) in complete DMEM supplemented with 2.5  $\mu$ g/ml 1,1'-dioctadecyl-3,3,3',3'-tetramethylindocarbocyanine perchlorate (DiI, Sigma-Aldrich, St. Louis, MO) for 30 minutes at 37°C, and subsequently washed three times with complete DMEM. Macrophages were incubated ( $1-10 \times 10^6$  cells/ml) in complete Macrophage medium, supplemented with 2.5  $\mu$ g/ml 3,3'-dioctadecyloxycarbocyanine perchlorate (DiO, Molecular Probes Inc, Paisley, UK) for 30 minutes at 37°C and subsequently washed three times with complete macrophage medium.

#### In vitro cytotoxicity assays

Cytotoxicity assays were performed by co-culturing DiI-labelled mouse B16F10 or B16F10-gp75 tumour cells and DiO-labelled macrophages in an effector to target (E:T) ratio of 15:1 in the presence of different concentrations of mouse anti-gp75 mIgG2a or human hIgG1, hIgG3, hIgG1-H435R or hIgG3-R435H. Human anti-HEPC IgG1 or mouse anti-Rat MG4 IgG2a antibodies served as isotype controls. Percentages of remaining tumour cells and macrophages that had taken up tumour cells (double-positive) were determined by flow cytometry after 24 hours of co-



culture. Percentages of tumour cells and double-positive macrophages after culture without mAbs were set at 100%.

#### Animal model

Mice were intraperitoneally (i.p.) injected with antibodies diluted in PBS (50µg/300µl) 4 days prior to injection of the tumour cells. B16F10-gp75 cells were harvested at the start of the experiment, washed and diluted in PBS to a concentration of 50.000 cells/300µl and i.p. injected. 14 days post injection mice were sacrificed with CO<sub>2</sub> and peritoneal metastasis were scored.

#### Statistical analysis

Data was analysed with Bonferroni-Post Hoc tests, preceded by two-way ANOVA tests for comparison of multiple groups. Significance was accepted at  $p < 0.05$ .


## **RESULTS AND DISCUSSION**

### **Antibody modification does not affect the binding capacity**

One potential drawback of using IgG3 isotype mAbs as therapeutic antibodies is the fact that human IgG1 (hIgG1) inhibits the recycling of hIgG3 via the neonatal FcRn, leading to a short half-life of IgG3<sup>10</sup>. Although FcRn binds both Fc domains of hIgG1 and hIgG3 with comparable affinity, the difference at position 435 in the Fc fragment of hIgG3 (histidine in IgG1 versus arginine in hIgG3) accounts for a decreased competitiveness of hIgG3 for pH-dependent FcRn binding and release, leading to increased hIgG3 lysosomal degradation<sup>10,21</sup>. To overcome the inferior half-life of wild-type hIgG3 we constructed a specific IgG3 containing a histidine at position 435 in its Fc domain, as it has been demonstrated that this leads to enhanced rescue through FcRn and subsequently to increased half-life *in vivo*<sup>10</sup>. In both the humanized hIgG1 TA99 as the hIgG3 TA99 mutations (histidine to arginine and vice versa respectively) were introduced at position 435 generating hIgG1-H435R TA99 and hIgG3-R435H TA99. Specific anti-human IgG1 or anti-human IgG3 ELISA confirmed the correct isotype of mAbs (Figure 1A and B). Additionally, we confirmed with page-gel electrophoresis that the structural integrity was not affected due to this amino acid modification (Supplementary Figure 1).

Next, the binding and functionality of human IgG1 and IgG3 TA99 variants was investigated. TA99 is directed against the 75-kD product of the *brown* gene, known as TRP-1 or glycoprotein gp75. This is highly conserved between mouse and human and expressed in melanocytic cells, like the mouse B16F10 melanoma cell line<sup>26</sup>. Gp75 is however exported efficiently towards intracellular melanosomes and almost no gp75 protein reaches the cell surface *in vitro*<sup>5,26-28</sup>. As such, mouse macrophages were not able to phagocytose B16F10 cells in the presence of TA99 *in vitro* (data not shown). It is possible that gp75 expression on B16F10 is up regulated *in vivo*, accounting for the *in vivo* efficacy seen with the anti-GP75 mIgG2a antibody<sup>29</sup>. This is supported by experiments performed with fresh B16F10 cells isolated from lung metastases, which had enhanced surface expression of gp75, and successful ADCC of these cells was induced through mIgG2a TA99 mAb *in vitro*<sup>30</sup>. Therefore, it is conceivable that *in vivo* TA99 mAbs are able to efficiently bind B16F10, and induce killing via high affinity interaction with murine Fcγ receptors. For *in vitro* studies, we therefore generated a cell line that stably expresses a truncated gp75 protein on the membrane by removing the dileucine intracellular retention signal motif which is critical for internalization (referred to as B16F10-gp75)<sup>28</sup>. TA99 hIgG1-H435R or hIgG3-R435H isoforms recognized membrane expressed gp75 on B16F10-gp75 cells

similarly (Figure 1C-D). In addition, titration of humanized IgG antibodies revealed similar binding curves, both between each other (Figure 1E) and compared to the original TA99 mouse IgG2a antibody (Figure 1F), indicating similar affinities.



**FIGURE 1. Generation of human IgG1 and IgG3 TA99 mAb with histidine-arginine rearrangements in Fc domains at amino acid position 435.**

Human IgG1 was mutated to contain an arginine at position 435 (IgG1 H435R), whereas the arginine at position 435 in human IgG3 was changed to histidine (IgG3 R435H). (A) Specific anti-human IgG1 or (B) anti-human IgG3 ELISA confirmed the correct isotype of mAbs. Staining of B16F10-gp75 with (C) different human TA99 mAb and (D) mouse TA99 IgG2a confirmed binding to surface gp75 and equal binding efficiency to gp75 of all human TA99 mAb. Concentration curves of human IgG1 and IgG3 mAb (E) and mouse IgG2a TA99 (F) on B16F10-gp75. Of note, scales of human (E) and mouse (F) antibodies are different.

anti-human IgG3 ELISA confirmed the correct isotype of mAbs. Staining of B16F10-gp75 with (C) different human TA99 mAb and (D) mouse TA99 IgG2a confirmed binding to surface gp75 and equal binding efficiency to gp75 of all human TA99 mAb. Concentration curves of human IgG1 and IgG3 mAb (E) and mouse IgG2a TA99 (F) on B16F10-gp75. Of note, scales of human (E) and mouse (F) antibodies are different.

### IgG1 is more efficient in inducing tumour cell kill in vitro with mouse macrophages

The efficiency of TA99 antibodies *in vitro* was tested by co-culturing mouse macrophages with the newly generated B16F10-gp75 cells in the presence of antibodies. At high antibody concentrations (2  $\mu\text{g}/\text{ml}$ ) hIgG1 and hIgG3 were equally efficient in inducing macrophage mediated tumour killing, with IgG1 mediating perhaps a slightly improved, but not significantly different, killing (Figure 2A).


Other studies demonstrated enhanced phagocytosis of red blood cells in the presence of hIgG3<sup>18,19,31-34</sup>. Moreover, opsonisation of *S. Typhimurium* with hIgG3 resulted in enhanced bacterial uptake by the monocyte cell line THP-1, which was dependent on Fc $\gamma$ RI expression<sup>35</sup>. We therefore tested whether TA99 hIgG3 was more efficient in B16F10-gp75 tumour cell killing and phagocytosis at lower, sub-optimal, mAb concentrations.

Lower concentrations of 0.1  $\mu\text{g}/\text{ml}$  hIgG1 or mIgG2a TA99 still induced B16F10-gp75 killing by mouse macrophages through phagocytosis (Figure 2B-D middle panel). Both hIgG1 and mIgG2a were equally able to induce tumour cell phagocytosis by macrophages (Figure 2C and D). By contrast, B16F10-gp75 killing was significantly reduced in the presence of human IgG3 TA99 mAb concentrations of 1  $\mu\text{g}/\text{ml}$  or lower (Figure 2C-D). *In vitro*, histidine or arginine rearrangements at amino acid position 435 in either hIgG1 or hIgG3 TA99 did not influence B16F10-gp75 killing (Figure 2C and D).

### Both human IgG1 and IgG3 inhibit formation of liver metastases

In other studies it has been demonstrated that transferring parts of the hIgG3 Fc domain into hIgG1 anti-CD20 mAb enhanced both complement binding, CDC and ADCC against lymphoma cells<sup>11,12,23</sup>. To study if TA99 hIgG3 has an improved effect

over TA99 hIgG1 and whether TA99 hIgG3-R435H overcomes the difference in half-life with IgG1 in an intra-peritoneal (i.p.) metastasis model of B16F10-gp75 tumour cells was used. Mice were injected i.p. with TA99 mAbs four days prior to i.p. injection of B16F10-gp75 and tumour growth was assessed 14 days post tumour cell injection. Conforming the *in vitro* data, mice treated with either mIgG2a or hIgG1 had almost no metastasis outgrowth (Figure 3). TA99 hIgG3, however, was less effective in preventing tumour outgrowth. Moreover, due to the introduced mutation in the TA99 hIgG1 recycling was expected to be reduced with as a result a decreased half-life. As a result, we hypothesised that this antibody would be less effective than the wildtype hIgG1. However, no difference between the wild type and the mutant was observed. Furthermore, mice treated with TA99 hIgG3 R435H were not better protected against tumour outgrowth compared to TA99 hIgG3 wildtype.




**FIGURE 2. Cytotoxicity assays using murine macrophages and B16F10-gp75 tumour cells.**

(A) Remaining B16F10-gp75 cells after a 24 hour incubation with macrophages and 2 µg/ml TA99 mAbs (different isotypes). (B) FACS analysis of co-cultures of DiO labelled murine macrophages (FL1) and DiI labelled B16F10-gp75 (FL2) tumour cells after 24 hours of treatment with 1 µg/ml mouse IgG2a or human IgG1 or IgG3 TA99 mAb. Macrophages, which have phagocytosed B16F10-gp75 tumour cells are encircled in FACS plots. (C) Percentage of remaining viable tumour cells and (D) increase in number of macrophages, which have phagocytosed B16F10-gp75 tumour cells after treatment of co-cultures with different concentrations of mAb. Mouse MG4 or human HEPC mAb were used as isotypes controls, which were set to 100%. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ , \*\*\*\* $P < 0.0001$ .

It has been demonstrated that mAb therapy of B16F10 liver metastases with mIgG2a TA99 (functional homologous to hIgG1) is independent of complement activation, but relies on Fcγ receptor and macrophage presence<sup>5,36</sup>. It has however not yet been investigated how hIgG3 TA99 exerts its effects. It is known that hIgG3 is superior in inducing complement subunit C1q binding compared to hIgG1<sup>17,37,38</sup>. Subsequently, C1q can activate the classical complement route, leading to complement induced lysis

of tumour cells. Additionally, immune responses can be induced by binding either directly to complement receptor 1 (CR1/CD35) on effector cells or via forming a complex with complement subunit 3, activating immune cells via complement receptor 3 (CR3/Mac-1/CD11b+CD18)<sup>39</sup>. CR1 is expressed by many different cells, including PMNs, monocytes, macrophages, and DCs but also B-cells and a subpopulation of CD4<sup>+</sup> T-cells<sup>40</sup>. Proliferation of B-and T cells is regulated by interaction of CR1 with C1q immune complexes, whereas PMNs, monocyte/macrophages and DCs are able to phagocytose complement opsonized immune complexes such as tumour cells. Mac-1/CR3 is primarily expressed on monocytes/macrophages and neutrophils, binds C3bi - the cleaved form of C3 - and contributes to ADCC. It was demonstrated that CD11b deficient mice that lack Mac-1 were less protected by treatment with mIgG2a TA99 mAb against B16F10 lung metastases development than wild-type mice<sup>41</sup>. Furthermore, although neutrophils were still able to penetrate malignant tissues and interacted with antibody opsonised tumour cells in Mac-1 <sup>-/-</sup> mice, they were unable to initiate ADCC towards tumour cells. Mac-1 on myeloid cells, including neutrophils therefore may play an important role in complement mediated tumour rejection. Because hIgG3 mAb were most effective in activating a respiratory burst in PMN, it is possible that the mechanism of action of hIgG3 in tumour rejection is dependent on both complement binding and PMN activation<sup>9,42</sup>. Mouse neutrophils however are not efficiently activated via IgG antibody. Therefore, mouse neutrophils are not likely to be effector cells in mAb-therapies tested in mouse models.



**FIGURE 3. Prevention of B16F10-gp75 metastases development with different TA99 isotype antibodies.**

Mice received peritoneal injections with 50µg mAb/mouse 4 days prior to peritoneal injection of B16F10-gp75 tumour cells. Metastases outgrowth after treatment with TA99 mouse IgG2a or (modified) human IgG1, human IgG3 mAbs was determined 14 days post tumour cell injection. \*\* $P < 0.01$ , \*\*\*\* $p < 0.0001$ .

Recently a study with chimeric anti-transferrin receptor 1 mAbs with an hIgG3 Fc domain demonstrated which amino acids are essential for both induction of ADCC and CDC<sup>43</sup>. When the leucines at position 234 and 235 were mutated into alanines (L234A, L235A) and the proline at position 331 into a serine (P331S) ADCC was significantly reduced. Mutating the P331S alone was sufficient to prevent CDC almost completely. Other studies demonstrated that IgG3 is superior in mediating complement activation and that the length of the hinge matters<sup>44,45</sup>. Additionally, correlations between the length of the hinge of IgG3 and activation of the complement at low antigenic-concentrations were made. This may enable hexamer-formation - required for efficient C1q binding - more efficiently at low epitope density<sup>46,47</sup>. This is supported by a recent report showing that CD20-specific IgG3 are indeed more efficient than IgG1 in mediating CDC against tumour targets - especially on those targets with low tumour antigen-expression such as chronic lymphocytic leukaemia (CLL). Using this

knowledge current therapeutic IgG1 antibodies might be enhanced.


A recent study has classified binding affinities of human IgG subclasses to mouse Fcγ receptors, and found human IgG3 to bind with superior affinity to all mouse Fcγ receptors except mouse FcγRI, to which IgG1 bound better<sup>48</sup>. Nevertheless, in our hands IgG1 was in most cases superior to that of IgG3 in mediating in *in vitro* and *in vivo* effector functions. A reason for this discrepancy has not yet been put forward, but may lay in the fact that increased binding of human IgG3 was also found for the inhibitory FcγRIIb, which may be expected to impede Fcγ receptor-mediated activities, without affecting complement-mediated functions.

In conclusion, humanized IgG3 does not show improved Fcγ receptor-dependent tumour cell killing with mouse effector cells *in vitro* and *in vivo*. Although literature suggests that IgG3 can have superior effector functions, this is likely mediated through activation of complement. Thus, although tumours that are susceptible for CDC may be killed more effectively via IgG3, Fc receptor-mediated functions will likely be less efficient. Overall, IgG3 mAbs may therefore be less suitable for antibody therapy of cancer.

## REFERENCES

1. Scott, A. M., Wolchok, J. D. & Old, L. J. Antibody therapy of cancer. *Nat. Rev. Cancer* **12**, 278–87 (2012).
2. Amoroso, A. *et al.* Understanding rituximab function and resistance: implications for tailored therapy. *Front. Biosci.* **16**, 770–82 (2011).
3. Braster, R., O’Toole, T. & van Egmond, M. Myeloid cells as effector cells for monoclonal antibody therapy of cancer. *Methods* **65**, 28–37 (2014).
4. Clynes, R., Takechi, Y., Moroi, Y., Houghton, A. & Ravetch, J. V. Fc receptors are required in passive and active immunity to melanoma. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 652–6 (1998).
5. Otten, M. A. *et al.* Experimental antibody therapy of liver metastases reveals functional redundancy between Fc gammaRI and Fc gammaRIV. *J. Immunol.* **181**, 6829–36 (2008).
6. van der Bij, G. J. *et al.* Experimentally induced liver metastases from colorectal cancer can be prevented by mononuclear phagocyte-mediated monoclonal antibody therapy. *J. Hepatol.* **53**, 677–85 (2010).
7. Gül, N. *et al.* Macrophages eliminate circulating tumor cells after monoclonal antibody therapy. *J. Clin. Invest.* **124**, 812–823 (2014).
8. Bruhns, P. *et al.* Specificity and affinity of human Fc gamma receptors and their polymorphic variants for human IgG subclasses. *Blood* **113**, 3716–25 (2009).
9. Vidarsson, G. *et al.* Activity of human IgG and IgA subclasses in immune defense against *Neisseria meningitidis* serogroup B. *J. Immunol.* **166**, 6250–6 (2001).
10. Stapleton, N. M. *et al.* Competition for FcRn-mediated transport gives rise to short half-life of human IgG3 and offers therapeutic potential. *Nat. Commun.* **2**, 599 (2011).
11. Rösner, T. *et al.* An IgG3 switch variant of rituximab mediates enhanced complement-dependent cytotoxicity against tumour cells with low CD20 expression levels. *Br. J. Haematol.* **161**, 282–6 (2013).
12. Natsume, A., Shimizu-Yokoyama, Y., Satoh, M., Shitara, K. & Niwa, R. Engineered anti-CD20 antibodies with enhanced complement-activating capacity mediate potent anti-lymphoma activity. *Cancer Sci.* **100**, 2411–8 (2009).
13. Vidarsson, G. & van de Winkel, J. G. Fc receptor and complement receptor-mediated phagocytosis in host defence. *Curr. Opin. Infect. Dis.* **11**, 271–8 (1998).
14. Bazin, R. *et al.* Use of hu-IgG-SCID mice to evaluate the in vivo stability of human monoclonal IgG antibodies. *J. Immunol. Methods* **172**, 209–217 (1994).
15. Sesarman, A., Vidarsson, G. & Sitaru, C. The neonatal Fc receptor as therapeutic target in IgG-mediated autoimmune diseases. *Cell. Mol. Life Sci.* **67**, 2533–50 (2010).
16. Carter, P. J. Potent antibody therapeutics by design. *Nat. Rev. Immunol.* **6**, 343–357 (2006).
17. Salfeld, J. G. Isotype selection in antibody engineering. *Nat. Biotechnol.* **25**, 1369–1372 (2007).
18. Wiener, E. *et al.* Differences between the activities of human monoclonal IgG1 and IgG3 subclasses of anti-D(Rh) antibody in their ability to mediate red cell-binding to macrophages. *Immunology* **62**, 401–404 (1987).
19. Tebo, A. E., Kremsner, P. G. & Luty, A. J. Fc gamma receptor-mediated phagocytosis of *Plasmodium falciparum*-infected erythrocytes in vitro. *Clin. Exp. Immunol.* **130**, 300–6 (2002).
20. Vidarsson, G., Dekkers, G. & Rispen, T. IgG Subclasses and Allotypes: From Structure to Effector Functions. *Front. Immunol.* **5**, 520 (2014).
21. Kim, J. K. *et al.* Mapping the site on human IgG for binding of the MHC class I-related receptor, FcRn. *Eur. J. Immunol.* **29**, 2819–25 (1999).
22. Einarsdottir, H. *et al.* H435-containing immunoglobulin G3 allotypes are transported efficiently across the human placenta: implications for alloantibody-mediated diseases of the newborn. *Transfusion* **54**, 665–71 (2014).
23. Natsume, A. *et al.* Engineered antibodies of IgG1/IgG3 mixed isotype with enhanced cytotoxic activities. *Cancer Res.* **68**, 3863–72 (2008).
24. Vidarsson, G., van de Winkel, J. G. J. & van Dijk, M. A. Multiplex screening for functionally rearranged immunoglobulin variable regions reveals expression of hybridoma-specific aberrant V-genes. *J. Immunol. Methods* **249**, 245–252 (2001).
25. Kimura, N., Du, R. P. & Mak, T. W. Rearrangement and organization of T cell receptor gamma chain genes in human leukemic T cell lines. *Eur. J. Immunol.* **17**, 1653–6 (1987).
26. Vijayasaradhi, S., Doskoch, P. M. & Houghton, A. N. Biosynthesis and intracellular movement of the melanosomal membrane glycoprotein gp75, the human b (brown) locus product. *Exp. Cell Res.* **196**, 233–240 (1991).
27. Zhou, B.-K., Boissy, R. E., Pifko-Hirst, S., Moran, D. J. & Orlow, S. J. Lysosome-Associated Membrane Protein-1 (LAMP-1) Is the Melanocyte Vesicular Membrane Glycoprotein Band II. *J. Invest. Dermatol.* **100**, 110–114 (1993).
28. Vijayasaradhi, S., Xu, Y., Bouchard, B. & Houghton, A. N. Intracellular sorting and targeting of

- melanosomal membrane proteins: identification of signals for sorting of the human brown locus protein, gp75. *J. Cell Biol.* **130**, 807–20 (1995).
29. Takechi, Y., Hara, I., Naftzger, C., Xu, Y. & Houghton, A. N. A melanosomal membrane protein is a cell surface target for melanoma therapy. *Clin. cancer Res.* **2**, 1837–42 (1996).
  30. Patel, D. *et al.* Enhanced suppression of melanoma tumor growth and metastasis by combined therapy with anti-VEGF receptor and anti-TYRP-1/gp75 monoclonal antibodies. *Anticancer Res.* **28**, 2679–86 (2008).
  31. Brouwers, H. A. *et al.* Maternal antibodies against fetal blood group antigens A or B: lytic activity of IgG subclasses in monocyte-driven cytotoxicity and correlation with ABO haemolytic disease of the newborn. *Br. J. Haematol.* **70**, 465–9 (1988).
  32. Rozsnyay, Z. *et al.* Distinctive role of IgG1 and IgG3 isotypes in Fc gamma R-mediated functions. *Immunology* **66**, 491–8 (1989).
  33. Perri, D., Shabani, F., Ziaeiian, G. & Denomme, G. A. IFN- $\gamma$  treated monocyte/macrophage phagocytosis of red cells sensitized with IgG1 and IgG3 Anti-D containing identical immunoglobulin variable region genes. *Transfus. Apher. Sci.* **39**, 37–44 (2008).
  34. Kumpel, B. M., De Haas, M., Koene, H. R., Van De Winkel, J. G. J. & Goodrick, M. J. Clearance of red cells by monoclonal IgG3 anti-D in vivo is affected by the VF polymorphism of Fc gamma RIIIa (CD16). *Clin. Exp. Immunol.* **132**, 81–6 (2003).
  35. Goh, Y. S. *et al.* Human IgG isotypes and activating Fc $\gamma$  receptors in the interaction of Salmonella enterica serovar Typhimurium with phagocytic cells. *Immunology* **133**, 74–83 (2011).
  36. Hara, I., Takechi, Y. & Houghton, A. N. Implicating a role for immune recognition of self in tumor rejection: passive immunization against the brown locus protein. *J. Exp. Med.* **182**, 1609–14 (1995).
  37. Recke, A. *et al.* Pathogenicity of IgG subclass autoantibodies to type VII collagen: induction of dermal-epidermal separation. *J. Autoimmun.* **34**, 435–44 (2010).
  38. Brüggemann, M. *et al.* Comparison of the effector functions of human immunoglobulins using a matched set of chimeric antibodies. *J. Exp. Med.* **166**, 1351–61 (1987).
  39. Bohlsón, S. S., O’Conner, S. D., Hulsebus, H. J., Ho, M.-M. & Fraser, D. A. Complement, C1q, and C1q-Related Molecules Regulate Macrophage Polarization. *Front. Immunol.* **5**, 402 (2014).
  40. Khera, R. & Das, N. Complement Receptor 1: Disease associations and therapeutic implications. *Mol. Immunol.* **46**, 761–772 (2009).
  41. van Spriel, A. B., van Ojik, H. H., Bakker, A., Jansen, M. J. H. & van de Winkel, J. G. J. Mac-1 (CD11b/CD18) is crucial for effective Fc receptor-mediated immunity to melanoma. *Blood* **101**, 253–8 (2003).
  42. AASE, A. & MICHAELSEN, T. E. Opsonophagocytic Activity Induced by Chimeric Antibodies of the Four Human IgG Subclasses With or Without Help from Complement. *Scand. J. Immunol.* **39**, 581–587 (1994).
  43. Leoh, L. S., Daniels-Wells, T. R., Martínez-Maza, O. & Penichet, M. L. Insights into the effector functions of human IgG3 in the context of an antibody targeting transferrin receptor 1. *Mol. Immunol.* **67**, 407–415 (2015).
  44. Michaelsen, T. E., Sandlie, I., Bratlie, D. B., Sandin, R. H. & Ihle, O. Structural difference in the complement activation site of human IgG1 and IgG3. *Scand. J. Immunol.* **70**, 553–64 (2009).
  45. Michaelsen, T. E. *et al.* Binding properties and anti-bacterial activities of V-region identical, human IgG and IgM antibodies, against group B Neisseria meningitidis. *Biochem. Soc. Trans.* **31**, 1032–5 (2003).
  46. Diebold, C. A. *et al.* Complement is activated by IgG hexamers assembled at the cell surface. *Science (80-. )*. **343**, 1260–3 (2014).
  47. Michaelsen, T. E., Garred, P. & Aase, a. Human IgG subclass pattern of inducing complement-mediated cytotoxicity depends on antigen concentration and to a lesser extent on epitope patchiness, antibody affinity and complement concentration. *Eur. J. Immunol.* **21**, 11–6 (1991).
  48. Overdijk, M. B. *et al.* Crosstalk between human IgG isotypes and murine effector cells. *J. Immunol.* **189**, 3430–8 (2012).

**SUPPLEMENTARY FIGURE 1****Structural integrity of hIgG1 and hIgG3 were not affected.**

Human IgG1 contains a histidine at position 435 to secure normal recycling via FcRn *in vivo*, whereas human IgG3 contains an arginine at this position, which hinders IgG3 recycling. Human IgG1 was mutated to contain an arginine at position 435 (IgG1 H435R), whereas the arginine at position 435 in human IgG3 was changed to histidine (IgG3 R435H). Non-reducing (left panel) and reducing (right panel) page-gel electrophoresis confirmed structural integrity of the different IgG1 and IgG3 mAbs.

