

# CHAPTER 4

## Diagnosis and management of acute appendicitis.

### EAES consensus development conference 2015

*Gorter RR, Eker HH, Gorter-Stam MA, Abis GS, Acharya A, Ankersmit M, Antoniou SA, Arolfo S, Babic B, Boni L, Bruntink M, van Dam DA, Defoort B, Deijen CL, DeLacy FB, Go PM, Harmsen AM, van den Helder RS, Iordache F, Ket JC, Muysoms FE, Ozmen MM, Papoulas M, Rhodes M, Straatman J, Tenhagen M, Turrado V, Vereczkei A, Vilallonga R, Deelder JD, Bonjer J.*

## **Abstract**

### **Background**

Unequivocal international guidelines regarding the diagnosis and management of patients with acute appendicitis are lacking. The aim of the consensus meeting 2015 of the EAES was to generate a European guideline based on best available evidence and expert opinions of a panel of EAES members.

### **Methods**

After a systematic review of the literature by an international group of surgical research fellows, an expert panel with extensive clinical experience in the management of appendicitis discussed statements and recommendations. Statements and recommendations with more than 70% agreement by the experts were selected for a web survey and the consensus meeting of the EAES in Bucharest in June 2015. EAES Members and attendees at the EAES meeting in Bucharest could vote on these statements and recommendations. In the case of more than 70% agreement, the statement or recommendation was defined as supported by the scientific community. Results from both the web survey and the consensus meeting in Bucharest are presented as percentages.

### **Results**

In total 46 statements and recommendations were selected for the web survey and consensus meeting. More than 232 members and attendees voted on them. In 41 of 46 statements and recommendations more than 70% agreement was reached. All 46 statements and recommendations are presented in this paper. They comprise topics regarding the diagnostic work-up, treatment indications, procedural aspects and postoperative care.

### **Conclusions**

The consensus meeting produced 46 statements and recommendations on the diagnostic work-up and management of appendicitis. The majority of the EAES members supported these statements. These consensus proceedings provide additional guidance to surgeons and surgical residents providing care to patients with appendicitis.

## Introduction

Acute appendicitis is a common gastrointestinal disease affecting 5.7 to 57 / per 100,000 individuals each year with the highest incidence in children and adolescents.<sup>1-6</sup> The variation of incidence is due to variations in ethnicity, sex, age, obesity and season of the year.<sup>3,6-</sup>

<sup>11</sup> Based upon the entrenched idea that appendicitis is an irreversible progressive disease eventually leading to perforation, removal of the appendix is the gold standard of treatment. The medical profession has gained much experience in managing patients with acute appendicitis ever since Fitz's first report in 1886.<sup>12</sup> Large heterogeneity exists, however, between existing intercontinental, European and national guidelines regarding diagnosing and managing acute appendicitis. For instance, in the Netherlands, preoperative imaging studies are promoted and considered mandatory in order to prevent negative appendectomies according to national guidelines, whereas in guidelines of other countries it is not promoted nor considered mandatory.<sup>13</sup> Another example is the inconsistency regarding the management of an unexpected "normal appendix" during diagnostic laparoscopy.<sup>13,14</sup> This heterogeneity prompted the need for an European consensus development conference for the diagnosis and management of acute appendicitis.

The European Association of Endoscopic Surgery (EAES) initiated a consensus development conference meeting on the management of acute appendicitis for its 2015 meeting in Bucharest. The aim of this consensus meeting was to develop practical guidelines based on the available evidence combined with the expertise of a selected panel of EAES surgeons. The findings are reported in this manuscript.

## Materials and Methods

The coordinating team (HJB, RG, HE and MGS) invited ten surgeons from nine European countries to serve as experts in this consensus development conference. An international research team of sixteen young surgical researchers across eleven European countries was formed to evaluate and process the existing literature on the management of acute appendicitis. The coordinators generated a list of topics on acute appendicitis to be addressed. (Appendix #1) An exploratory literature search was conducted in order to identify any additional topics of interest. All topics were approved by the experts and subsequently divided into three main parts: preoperative care, operative care and after care. Based upon the topics, research questions were formulated, reviewed and approved by the panel of experts.

### Literature Search and processing of the Literature

Research questions were used as guidance to conduct literature searches. The searches were conducted in cooperation with a medical information specialist of the Vrije Universiteit. Searches were performed in the following databases: Pubmed, Web of science and the Cochrane library from inception up to 31<sup>st</sup> of December 2014. No limitation was used regarding year of publication. Searches have been attached in Appendix #2. All papers published in European languages and all study types with the exception of casereports were included in the search.

All articles were screened and reviewed by teams of two research fellows for eligibility, based on title and abstract. If eligible for inclusion, full text articles were obtained. If no full text was available, the article was excluded. In case of disagreement between the two research fellows, the coordinator dedicated to the topic acted as referee. Full text articles were summarised, evaluated and discussed at research meetings to assess their eligibility for inclusion in the review process. All included studies were evaluated according to the GRADE system.<sup>15-18</sup> The GRADE system systematically evaluates the available literature and focusses on the level of evidence based upon the types of studies included. The level of evidence can be marked as high, moderate, low or very low. This could either be downgraded in case of significant bias or upgraded when multiple high-quality studies showed consistent results. The highest levels of evidence (systematic reviews) were assessed first. If the systematic review was of sufficient quality, it was used to answer the research question. If no systematic review of sufficient quality was found, randomised controlled trials (RCTs) and cohort studies were evaluated. All selected studies were uploaded to a Mendeley database that was accessible to all research fellows, coordinators and experts.

After the literature search, an expert was assigned to every couple of researchers. This threesome was assigned research questions from the preoperative care, operative care and after care. Hereafter they were responsible for formulating a statement/conclusion and, if possible, a recommendation on the assigned research questions. Again, the quality of the evidence was evaluated according to the GRADE/SIGN system.<sup>15-19</sup> The strength of the recommendation was based on the level of evidence and qualified as weak or strong. This was reflected in terms, using “*recommend*” in case of a strong recommendation and “*suggest*” in case of a weak recommendation.

A face-to-face consensus meeting among the experts was held in Amsterdam on the 1<sup>st</sup> of May 2015 to discuss the final statements and recommendations. The coordinating team, all experts and members of the international research team attended the meeting. A modified Delphi method was used. The Delphi method is a structured process, commonly used to develop healthcare quality indicator and consists of four key components; iteration, controlled acquisition of feedback, aggregation of responses and anonymity. As anonymity was not applicable in our situation, we used the term modified.<sup>20-22</sup> All statements and recommendations were shared with proposed level of evidence with the entire group. After displaying the statements and recommendations, the experts casted their votes of agreement or disagreement. Refrain from voting was not allowed. No discussion was allowed between the experts at this point of time. In case of 100% consensus, the statement and recommendation were accepted without further voting or discussion. In case of lack of consensus, the research team responsible for the statement presented the underlying considerations. After discussion between the experts, a second voting round was conducted. The statement or recommendation was accepted in case of at least 70% consensus. Those statements and recommendations with less than 70% consensus in the expert meeting were not included in the web survey or in the 2015 Bucharest meeting.


All finalised recommendations and statements with levels of evidence were entered into a web survey and distributed to all EAES members by e-mail. The web survey was open from May 27<sup>th</sup> until July 3<sup>rd</sup> 2015. The recommendations or statements as well as the levels of evidence were open to several voting options: ‘agree’, ‘partly agree’, ‘disagree’ or ‘don’t know’. The option “partly agree” meant that the voter agreed with the recommendation, but did not agree with the strength of recommendation.

All finalised recommendations and statements from the Amsterdam meeting with levels of evidence were presented at a plenary session of the 23<sup>rd</sup> annual meeting of the EAES on the 5<sup>th</sup> of June 2015 in Bucharest. Live voting was performed using a digital voting system. Voting options were the same as the abovementioned.

Both results from the web survey and the Bucharest meeting are presented in the results-section.

## Results

The literature search yielded 13,132 articles. The title, abstract and full text were reviewed. In total, 675 articles were selected and reviewed in detail to define 75 statements and recommendations, which were subsequently discussed at the Amsterdam meeting. (Appendix #1) During this meeting, the following statements and recommendations were excluded: on incidence and prevalence of appendicitis (n=4), on the place of NOTES in acute appendicitis (n=1), on the learning curve of appendectomy (n=1), on day surgery for acute appendicitis (n=1) and on the skeletonizing technique of the mesoappendix (n=1). Twenty-one statements were combined leaving a total of 46 statements and recommendations; 8 statements and 14 recommendations for preoperative care, 1 statement and 15 recommendations for operative care and 2 statements and 6 recommendations for aftercare. (Figure #1) Of the 675 articles, 100 were excluded due to the fact that statements and recommendations were excluded or were combined, rendering 575 articles. (Figure #1 / Appendix #3)



**Figure 1.** Flow diagram of the process prior to the EAES consensus meeting in Bucharest 2015.

**Web survey**

In total, 317 EAES members responded to the web survey, 90% were surgeons and 10% surgical residents.

**Bucharest meeting**

The 2015 EAES congress in Bucharest was attended by 1166 delegates. During the plenary consensus meeting, 232 delegates voted. Sixty-eight percent were surgeons, 26% surgical residents and 6% scientists, physician assistants and others.

**Diagnosis (Table 1-4)****Preoperative care**

Establishing the diagnosis of acute appendicitis remains challenging. The clinical presentation of acute appendicitis can vary from mild symptoms to signs of generalised peritonitis and sepsis. Hence, the value of individual clinical variables to determine the likelihood of acute appendicitis in a patient is low.<sup>23,24</sup> Biochemical testing is performed routinely in most patients. Its value in confirming acute appendicitis is debatable. A recent systematic review showed that elevated C-reactive protein levels render the highest diagnostic accuracy followed by increased numbers of leucocytes with an area under the curve of 0.75 [95%CI: 0.71-0.78] and 0.72 [95%CI: 0.68-0.76] respectively.<sup>24</sup> The area under the curve represents the ability of a test to correctly classify patients. In case the score is between the 0.7 and 0.8, it represents a fair test. Both clinical and biochemical variables have been combined into clinical predicting rules (CPR) such as the Alvarado score and paediatric appendicitis score (PAS).<sup>25,26</sup> This was done to increase the value of the individual variables. Ohle et al demonstrated that the Alvarado score was good at “ruling-out” appendicitis with an overall sensitivity and specificity of 96% and 81% respectively.<sup>27</sup> In children, however, it has been shown that the PAS outperforms the Alvarado score.<sup>28</sup> To increase the predictive value of these two tests Ebell et al identified new cut-off values for the Alvarado score and PAS, which improved sensitivity and specificity rates.<sup>29</sup> Based upon the Alvarado score, patients can now be categorised into low risk (Score < 4), intermediate (4-8) and high risk ( $\geq 9$ ). The use of such CPRs appears useful to determine the likelihood of acute appendicitis. Distinguishing between low, intermediate and high risk provides guidance whether imaging studies are necessary.


Imaging studies in patients with a clinical suspicion of acute appendicitis can reduce the negative appendectomy rate, which has been reported to be as high as 15%. Ultrasonography, abdominal computed tomography (CT) and magnetic resonance imaging (MRI) are most commonly used. Ultrasonography is non-invasive, avoids radiation and is associated with a sensitivity rate between 71 and 94% and a specificity rate between 81 and 98%. The positive likelihood ratio of ultrasonography is high at values between 6 and 46 while the negative likelihood ratio is moderate (0.08 - 0.30).<sup>30-39</sup> Ultrasonography is therefore reliable to confirm

presence of appendicitis but unreliable to exclude appendicitis. Furthermore, one should bear in mind that ultrasonography is highly operator dependent. Inconclusive ultrasonography findings, mostly due to failure visualising the appendix, mandate additional imaging studies.

Abdominal computed tomography (CT) for suspected appendicitis has sensitivity and specificity rates between 76-100% and 83-100% respectively and, therefore, is superior to ultrasonography. Lower values of sensitivity and specificity can be explained by the use of enteral contrast.<sup>32,33,35-44</sup> However, the radiation exposure of abdominal CT is a concern particularly in children and during pregnancy. The estimated lifetime cancer related mortality risk of developing a radiation-induced malignancy is approximately 0.18% for a one-year-old child and 0.11% in a 15-year-old child if an abdominal CT is performed.<sup>45,46</sup> Computed tomographies employing only a quarter of the standard radiation dose (low-dose CTs) provide similar imaging results as standard CTs and are, hence, an excellent alternative.<sup>47</sup> Regarding the administration of oral contrast, Andersson et al concluded in their meta-analysis that a CT-scan without oral contrast was superior to CTs with oral contrast in terms of sensitivity and specificity.<sup>48</sup> Therefore, low-dose CTs without oral contrast are preferable in patients with suspected appendicitis.<sup>48</sup>

Magnetic resonance imaging (MRI) is used in pregnant patients and children with inconclusive findings at ultrasonography.<sup>49</sup> A recent meta-analysis on MRI in 363 patients with appendicitis, yielded a sensitivity rate of 97% [95%CI: 92-99%], a specificity rate of 95% [95%CI: 94-99%], a positive likelihood ratio of 16.3 [95%CI: 9.10-29.10] and a negative likelihood ratio of 0.09 [95%CI:0.04-0.20].<sup>50</sup> These rates are comparable to those of CT-imaging, although these findings should be interpreted with care as most studies have been performed in a selected group of patients. MRI is associated with significant costs and interpreting the images requires experience. Therefore, at the present time, use of MRI appears limited to pregnant women and children.

The algorithm associated with the Alvarado score (recommendation 4) is shown in figure 2.



**Figure 2.** Algorithm

- \* The cut-off values are based upon the study by Ebell et al<sup>29</sup>
- \*\* One could consider performing additional imaging studies in patients with high probability based upon the Alvarado score in order to reduce the negative appendectomy rate.
- \*\*\* Ultrasound should be performed as a first level diagnostic imaging study, although in specific patient groups (such as the obese) an immediate CT might be considered.
- \*\*\*\* In case of an inconclusive result from the ultrasound, we recommend that additional imaging studies should be performed. Either a CT or MRI is preferred although it is recommended to perform a MRI in children and pregnant patients. It is therefore obligated to rule out pregnancy before a CT is obtained in a woman of reproductive age suspected of appendicitis
- \*\*\*\*\* In case all the imaging studies are inconclusive, patients should be observed and reassessed. Diagnostic laparoscopy should be reserved for those patients with a continuous high index of suspicion after reassessment.
- \*\*\*\*\* In case of low probability based upon the Alvarado score, other diagnoses should be excluded and the patient can either be discharged with good instruction (with an optional reassessment the next day) or admitted for observation if the clinical condition mandates this. In case appendicitis is excluded, patients should be treated for the set diagnosis according to the local protocols.

In obese patients (definition depends on the reference study), the diagnostic accuracy of ultrasound is diminished due to an increase of the subcutaneous and intra-abdominal fat. Anderson et al demonstrated that the body mass index (BMI) does not alter the diagnostic accuracy of a CT.<sup>51</sup> CT appears therefore more reliable than ultrasonography in obese patients with the exception of children and pregnancy.

### **Treatment indications (Table 1-4)**

Patients with appendicitis are classified as uncomplicated or complicated appendicitis based upon preoperative, intraoperative and/or histopathological findings. In this report, uncomplicated appendicitis has been defined as an inflamed appendix without signs of gangrene, perforation, intraperitoneal purulent fluid, contained phlegmon or intra-abdominal abscess (IAA). Complicated appendicitis applies to all patients with either a gangrenous inflamed appendix with or without perforation, intra-abdominal abscess, periappendicular contained phlegmon or purulent free fluid. Classification is necessary as treatment strategies may differ.

#### *Uncomplicated appendicitis*

Appendectomy is still considered to be the gold standard for uncomplicated appendicitis. Two main approaches to remove an inflamed appendix are available; the open approach (OA) or the laparoscopic approach (LA). In 2010 a large Cochrane review on 67 studies showed that LA significantly reduced the rate of surgical site infection (SSI) (OR 0.43 [95%CI: 0.34-0.54]) but significantly increased the risk of an intra-abdominal abscess (IAA) (OR 1.77 [95%CI: 1.14-2.76]) compared to the open approach.<sup>52</sup> It was stated that LA was associated with fewer superficial wound infections, less postoperative pain, shorter hospital stay and earlier return to work, but the higher rate of IAA raised concerns.<sup>52</sup> Ever since, inconsistent results have been reported regarding the potential higher incidence of IAA after LA.<sup>53-61</sup> Benefits of LA over OA reported in meta-analyses are: reduced incidence of SSI, postoperative and long-term bowel obstruction with better outcome in terms of shorter hospital stay, its diagnostic value, less pain, earlier return to work, earlier start of oral intake, improved scar and body satisfaction and fewer incisional hernias.<sup>54,55,58,61-66</sup> Disadvantages beside the possible higher incidence of IAA are longer operative time and possibly increased costs.<sup>58,63</sup>

To reduce the surgical trauma even more, new treatment strategies have been introduced such as Single Incision Laparoscopic Surgery (SILS) first reported by Pelosi et al.<sup>67</sup> Since then, numerous studies (RCTs and SR) have been published on the potential advantages and disadvantages of the SILS technique. It can be concluded that SILS is associated with comparable postoperative morbidity rates compared to conventional LA.<sup>68-70</sup> The disadvantage is the fact that SILS is a more difficult technique as is reflected by the higher technical failure rate, longer operative time and conversion rate.<sup>71-78</sup> Main advantages of SILS would be less postoperative pain and better cosmetic outcomes, although inconsistent results have been reported.<sup>71,75,76,79-81</sup> At the present time, evidence is lacking that SILS is superior to conventional LA.<sup>79,82,83</sup> SILS is, however, a safe and feasible alternative.

Recently, initial non-operative management of appendicitis has been investigated in the adult population. Five RCTs reported an effectiveness of 41-85% at one-year follow up.<sup>84-88</sup> Meta-analyses of these studies revealed that non-operative treatment of acute appendicitis is less

effective but could avoid surgery in 60-85% of patients.<sup>89-94</sup> Opponents of this strategy raise concerns such as recurrent appendicitis, missing an underlying malignancy and progression of uncomplicated into complicated appendicitis. Due to the possible avoidance of surgery with an initial non-operative treatment strategy, morbidity was diminished.<sup>91,93,95</sup> However, both RCTs and meta-analyses show significant heterogeneity of methodological quality, studies included and definitions of outcome parameters. Until higher qualitative evidence has been obtained regarding the potential benefits of initial non-operative management of acute appendicitis and the potential long-term effects have been investigated appropriately, appendectomy remains the gold standard in acute uncomplicated appendicitis.

### *Complicated appendicitis*

Due to the heterogeneity of the definitions used in the literature, it is difficult to draw firm conclusions regarding the treatment of complicated appendicitis. In 2013 Dimitriou et al published a retrospective cohort study on 150 patients with complicated appendicitis (defined as perforated appendicitis with an abscess or peritonitis). They showed that LA reduced the incidence of SSI, number of reoperations and length of hospital stay as compared to OA with no difference in IAA rate.<sup>96</sup> An RCT encompassing 81 patients with clinically and histopathologically confirmed complicated appendicitis showed similar outcomes after OA and LA.<sup>97</sup> It should be noted however, that the incidence of IAA after LA for patients with complicated appendicitis was reported to be higher in some studies. Tuggle and colleagues reported that LA in patients with complicated appendicitis was associated with an incidence of IAA of 6.7% versus 3.7% in patients who underwent an open appendectomy.<sup>98</sup> The incidence of small bowel obstructions after LA is lower compared to OA (pooled odds ratio 0.44 [95%CI: 0.26-0.74] with large heterogeneity regarding follow-up period).<sup>65</sup>

In case of a contained phlegmon or abscess (periappendicular mass) some authors opt for non-operative treatment while others advocate aggressive operative treatment. In 2007 Andersson et al demonstrated that immediate surgical treatment of patient with an abscess or phlegmon was associated with higher morbidity compared to initial non-operative treatment (Odds ratio: 3.3 [95% CI: 1.9-5.6]).<sup>99</sup> Similis et al showed in their meta-analysis of 17 studies regarding this specific patient group that non-operative treatment was associated with fewer complications (SSI, IAA and bowel obstructions). It must be mentioned that this meta-analysis was subject to large heterogeneity.<sup>100</sup> Recent cohort studies draw opposite conclusions.<sup>101,102</sup> They opt for a more aggressive surgical approach at time of presentation in case of an appendicular mass or appendicular abscess, based upon the idea that there is a relative high failure rate for non-operative treatment.<sup>101,102</sup> In our opinion, with this new evidence, a new systematic review should be performed. Until then, initial non-operative treatment of an appendicular mass or appendicular abscess is the preferred treatment of choice. Although not covered in this consensus guideline, the value of interval appendectomy after initial non-

operative treatment of an appendicular mass is still subject of debate. Some opt for an interval appendectomy based upon the chance of missing an underlying and untreated malignancy (incidence 6%) and the chance of developing recurrent appendicitis (incidence 5-44%).<sup>101-103</sup> Both can be avoided with an interval appendectomy, although data are lacking on its benefits.

#### *Specific patient groups*

- Obese patients

Abdominal surgery in obese patients is challenging for both the anaesthesiologist and surgeon due to higher incidence of respiratory dysfunction, difficult access to the abdominal cavity, blurred anatomical landmarks and reduced working space in the abdominal cavity. Clarke et al performed a subgroup analysis among 37 patients (14 LA and 23 OA) with a BMI higher than 30 kg/m<sup>2</sup>.<sup>104</sup> and reported similar morbidity after LA and OA.<sup>104</sup> This was confirmed by a meta-analysis, although a reduced length of hospital stay was noted after LA.<sup>105</sup> More recently, two recent meta-analyses showed a reduction of mortality and morbidity rates after LA.<sup>106,107</sup>

- Pregnancy

Pregnancy induces anatomical and physiological changes that challenge the surgeon. The potential effects of carbon dioxide and increased abdominal pressure during LA on the foetus remain unclear. Loss of the foetus is most feared. In 2008 Walsh et al published a systematic review of 637 laparoscopic appendectomies in pregnant patients and noted foetal loss in approximately 6% of the patients, with the highest incidence in patients with complicated appendicitis.<sup>108</sup> Another review confirmed these findings and reported a nearly twofold increase of foetal loss in the LA group.<sup>109</sup> Both reviews, however, are mainly dominated by one study and based on low-grade evidence (retrospective studies with small numbers of patients).<sup>108-110</sup> Recently, a review suggested that based upon the little available evidence no recommendation can be made regarding the preferred approach in pregnant patients.<sup>111</sup> More studies are necessary to ascertain the role of laparoscopic surgery during pregnancy. Until more evidence comes available, the surgical approach should be at the surgeon's discretion. Based upon expert opinion, we recommend laparoscopy in case of sufficient experience. Although not supported by literature, we strongly advise a multi-disciplinary approach to the pregnant patient with appendicitis.<sup>13,54,82,111,112</sup>


- Children

One meta-analysis included 107,624 children with both uncomplicated and complicated appendicitis.<sup>113</sup> Laparoscopic appendectomy in children with uncomplicated appendicitis was associated with a significant reduction of hospital stay with similar morbidity compared to open surgery. In children with complicated appendicitis, LA was associated with lower rates of morbidity, SSI, length of hospital

admission and bowel obstruction. However, laparoscopic surgeries lasted longer and were followed by more intra-abdominal abscesses.<sup>113</sup> In more recent prospective cohort studies in children below 5 years of age, LA was associated with fewer complications.<sup>114</sup> Non-operative treatment of acute uncomplicated appendicitis appears more promising in children than in adults.<sup>115,116</sup>


- Elderly  
Elderly patients have higher morbidity, reduced physiological reserves and impaired inflammatory responses, which increases their perioperative risks. All studies of laparoscopic appendectomy in elderly support the use of laparoscopic surgery.<sup>117-121</sup> One meta-analysis, comprising more than 15,000 patients reported that LA reduced postoperative mortality (0.24 [95%CI: 0.15-0.37]), postoperative complications (0.61 [95%CI: 0.50 to 0.73]) and length of hospital stay (-0.51 [95%CI: -0.64 to - 0.37]) compared to OA.<sup>119</sup>

**Table #1.** Preoperative care: Statements EAES meeting 2015




\* Abbreviation: LOE = Level of evidence

**Table #2.** Preoperative care: Statements Web survey




\* Abbreviation: LOE = Level of evidence

**Table #3.** Preoperative care: Recommendations EAES meeting 2015



\* Abbreviation: LOE = Level of evidence, SOR = Strength of Recommendation

**Table #4.** Preoperative care: Recommendations Web survey



\* Abbreviation: LOE = Level of evidence, SOR = Strength of Recommendation

**Operative care (Table 5-8)****Timing**

Determining the best moment to perform surgery in case of acute appendicitis is of crucial importance.<sup>122,123</sup> Acute appendicitis has been considered to be an irreversible progressive disease although recent studies have questioned this dogma.<sup>84,89,124</sup> Nowadays, the idea is endorsed that two types of appendicitis exist: uncomplicated (nonperforating) and complicated (perforating) appendicitis. The aetiology and pathogenesis of acute appendicitis remain largely unknown. Predicting a mild or fulminant course of appendicitis is not possible. Delaying an appendectomy increases the risk of perforated appendicitis, which is associated with higher incidence of short- and long-term morbidity.<sup>125-127</sup> Hence, it is recommended to perform appendectomy as soon as possible. Although it should be noted that some studies have revealed that the clinical outcome was not affected by time to surgery (when this was performed within 12 hours after presentation at the emergency department).<sup>128-129</sup>

**Antibiotic prophylaxis**

Antibiotic prophylaxis has been proven effective in prevention of superficial surgical site infections and intra-abdominal abscesses in patients with appendicitis.<sup>130-132</sup> Prophylaxis should be commenced at the time of establishing the diagnosis of acute appendicitis. The choice of antibiotics is dependent on the local microbiome and drug resistance pattern and is not influenced by age.

**Technique**

Open access to the abdominal cavity as well as closed access using the Veres needle are accepted techniques to perform laparoscopy.<sup>133-135</sup> The debate on the preferred technique continues. However, in children, the majority of surgeons employs open establishment of a pneumoperitoneum.

The placement of the camera-port and the work-ports depend on the anatomy of the patient and preference of the operating surgeon. Primary principle of trocar placement in laparoscopy is that a triangular working space should be pursued.

**Intraoperative procedure**

Increased employment of preoperative radiologic testing (e.g. ultrasound, CT or MRI) in cases of suspected appendicitis has significantly reduced the incidence of a normal appearing appendix encountered during surgery.<sup>136</sup> Macroscopic distinction between a normal appendix and appendicitis during surgery can be difficult.<sup>137,138</sup> The “gold standard” for defining appendicitis is histopathology. In some studies, histopathological assessment revealed abnormal findings in up to 26% of macroscopically normal appearing appendices.<sup>139,140</sup> Therefore, it is recommended to perform an appendectomy in case of a normal appearing appendix during surgery for suspected appendicitis.

Several studies have investigated the safety of different methods of securing the appendicular stump.<sup>82,141-143</sup> None of the different closure methods has a clear advantage in case of a healthy appendix base. Stapler devices provide the most standardised and patent closure of the appendix base. Suturing of the appendix base provides sufficient closure as well, but is technically more demanding than other techniques.<sup>142</sup> In case of perforation of the appendicular base, clips or endoloops do not provide secure closure and staple devices or laparoscopic suturing are required.<sup>82</sup>


Reduction of bacterial load by meticulous suction of intraperitoneal fluids is advised.<sup>144-146</sup> The right paracolic and pelvic area should be inspected to leave no fluid collections behind. Irrigation of the intraperitoneal space in case of perforated appendicitis seems to be counterproductive leading to a higher number of abscesses.<sup>144-145</sup> It is believed that irrigation of the intraperitoneal space leads to spreading of bacteria. Routine use of drains does not reduce the incidence of abscesses.<sup>145,147</sup> Necessity of a drain for special indications is left to the discretion of the surgeon.

### **Intraoperative unexpected findings**

When an appendicular mass is encountered during surgery one should restrain from continuing the operation. Continuation of the operation can necessitate bowel resection. Antibiotic treatment of phlegmon and drainage of any abscess should be performed.<sup>99,148,149</sup>


The extent of surgical resection in case of suspected malignancy depends on the location and size of the appendicular mass.<sup>150-154</sup> Routine inclusion of the mesoappendix with the appendectomy is advised. Definitive histological findings determine if an additional resection after appendectomy is indicated. In cases of small neuro endocrine tumours (NET) or low-grade appendicular mucinous neoplasms (LAMN), a total mesoappendicular resection can be sufficient. In cases of a NET >1cm, LAMN grade 3-4 or an adenocarcinoma of the appendix a formal right hemicolectomy is indicated to provide an oncologically sufficient resection. It is advised to perform a total meso-appendicular resection at the primary operation and an additional hemicolectomy at a later stage when indicated.<sup>150-154</sup>

**Table #5.** Operative care: Statements EAES meeting 2015




\* Abbreviation: LOE = Level of evidence

**Table #6.** Operative care: Statements Web survey




\* Abbreviation: LOE = Level of evidence

**Table #7. Operative care: Recommendations EAES meeting 2015**



\* Abbreviation: LOE = Level of evidence, SOR = Strength of Recommendation

**Table #8.** Operative care: Recommendations Web survey



\* Abbreviation: LOE = Level of evidence, SOR = Strength of Recommendation

## **Aftercare (Table 9-12)**

### **Postoperative antibiotics**

The incidence of SSI after appendectomy has been reported to range from 0-11%.<sup>155-164</sup> The severity of appendicitis strongly influences the risk of developing postoperative complications resulting in a substantially higher complication rate (up to 2-4 times) in patients with complicated appendicitis. In this specific group, postoperative administration of antibiotics significantly reduces the rate of SSI. In addition, to reduce bacteraemia and sepsis, these patients are uniformly treated with a course of postoperative antibiotics.<sup>155-158,163</sup> In uncomplicated appendicitis there is no evidence supporting routine administration of postoperative antibiotics. Therefore, only one preoperative dose is advised.<sup>155-158</sup>

Advice on type of antibiotics depend on local microbiome and resistance patterns, and therefore should be left up to the discretion of the surgeon.<sup>159,160</sup> Available evidence on duration of treatment is limited and mainly focussed on children. However, there is no firm evidence on the duration (3, 5, 7, 10 days) and route of administration (usually initial intravenous administration for 48 hours, then oral administration).<sup>156,157,159,161,162</sup>

### **Postoperative complications**

The incidence of postoperative complications ranges from 3.0 to 28.7%.<sup>164-174</sup> Complications include small bowel obstruction (0 – 1.9%), SSI (1.2 – 12.0%), IAA (1.6 – 8.0%), stump leakage and stump appendicitis.<sup>164-174</sup> Literature suggests a higher rate of complications in complicated appendicitis.<sup>166,167,171,175</sup>

Literature on stump leakage and stump appendicitis is limited, no exact incidences have been reported in the literature, although it is assumed that it is more common in patients with complicated appendicitis and after OA.<sup>176</sup> A recommendation to avoid stump leakage or stump appendicitis is to resect the appendix as a whole.<sup>176</sup> Therefore, the stump should be no longer than 0.5 cm and caecal tenia should be followed onto the appendix at removal to ensure complete resection. Stump appendicitis is significantly more associated with perforation, as diagnosis is delayed by misled attention. This is caused by the assumption that the appendix as a whole is resected. Prevention is crucial. In case of timely diagnosis stump resection with laparoscopic or open approach is feasible. In case of perforation usually extended bowel resection is required.<sup>176</sup>

In the initial management of IAA after appendectomy conservative measures (i.e. non-operative with antibiotics) are effective in most patients. However, in case of lack of improvement or deterioration, a more invasive strategy should be applied (percutaneous drainage or surgical (laparoscopic) drainage).<sup>177-179</sup>


**Postoperative care**

The use of prophylactic anti-emetics diminishes the incidence of post-operative nausea and vomiting. Increasing the diet is best determined by the patient’s ability to tolerate oral intake. There is no evidence that a liberal diet causes complications in the postoperative period.<sup>164,180</sup> Postoperative pain management should follow local protocol for pain management after abdominal surgery. Postoperative analgesia with PCA provides effective and safe pain relief in children and adults and is less time-costly.<sup>181</sup> Recently positive results have been published regarding the pre-emptive incision site infiltration with a local anaesthetic. Studies demonstrated that this decreases the total opioid consumption and lowers pain score experienced by patients in the first 24 hours after surgery.<sup>182-184</sup>

**Pathology**


Carcinoid is the most commonly found neoplasm in appendectomy specimens with an incidence between 0.13-2.4%.<sup>185-190</sup> Other unexpected findings can be countered in 1.4-2.4% of patients, including: diverticulitis (1.2%), tuberculosis appendix (0.1%), endometriosis (3.6%), adenocarcinoma (<1%) and mucinous cystadenoma (0.2-0.6%).<sup>191-194</sup> Treatment of unexpected findings ranges from no further surgical treatment, to right hemi colectomy and even hyperthermic intraperitoneal chemotherapy (HIPEC) in some cases.<sup>195-197</sup> Even though the incidence of unexpected findings seems low, the actual number of patients is significant and correct diagnosis is crucial for adequate treatment.<sup>198-202</sup>

**Table #9.** After care: Statements EAES meeting 2015




\* Abbreviation: LOE = Level of evidence

**Table #10.** After care: Statements Web survey




\* Abbreviation: LOE = Level of evidence

**Table #11.** After care: Recommendations EAES meeting



\* Abbreviation: LOE = Level of evidence, SOR = Strength of Recommendation

**Table #12.** After care: Recommendations Web survey



\* Abbreviation: LOE = Level of evidence, SOR = Strength of Recommendation

## Discussion

This EAES consensus development conference regarding the diagnosis and management of acute appendicitis resulted in 46 statements and recommendations based upon the available evidence. Results from this meeting led to this paper, which can be used as a guideline for surgeons treating patients with appendicitis. Local guidelines, national guidelines and guidelines from scientific communities regarding appendicitis were available but showed great heterogeneity.<sup>13,14,203</sup> With this consensus meeting we managed to gather experts from different European nations to compare and debate management of patients with acute appendicitis. This led to a consensus meeting in which 41 of the 46 statements and recommendations were supported by the majority of the members of the EAES. The transfer of knowledge between the member countries, the opportunity to discuss views and above all the creation of a widely supported paper appears valuable.

Our list of topics was created by the coordinating team and expert panel and was thought to cover the most important topics in the field of acute appendicitis. Despite local differences, the general idea within the consensus group on the management of patients with acute

appendicitis was comparable. In some cases differences of treatment strategies between members of the expert panel were due to available surgical supplies and finances. This is reflected for instance on the statements and recommendations regarding SILS and MRI. However, we want to emphasize that in defining statements we refrained from stating specific procedures. We rather stated the general principles to follow. In this way, the results from this consensus guideline can also be applied in areas with limited resources.

The methodology of a consensus guideline is always subject to discussion. In the literature there are several ways to conduct consensus conferences.<sup>20,204-206</sup> However, not one was suited for our situation. It was therefore decided to modify the Delphi method, as described in the method section, in order to systematically evaluate each statement and recommendation.<sup>20-22</sup> We decided to finalise only those statements and recommendations with 70% or more consensus, which is the arbitrary cut-off value we selected. The results of both the websurvey and the live voting at the EAES conference in Bucharest are presented independently rather than combined to rule out any bias. As expected, small differences were noted between the several voting rounds. Although supported by the experts, some statements and recommendations were not supported by the scientific community in both the web survey as in the Bucharest meeting. The topics that were not supported were on accuracy of MRI compared to CT, the application of SILS, extensive work-up in the elderly and treatment strategy for immune compromised patients and the open access to the peritoneal cavity. Explanations for these discrepancies might be related to local habits, experience and financial situation. Of more interest are the discrepancies noted between the outcome in the web survey and during the Bucharest meeting. Discrepancies were noted on the topic of MRI application in children, the preferred approach in pregnant patients and the use of local anaesthetics prior to incision. This can again be explained by the fact that local habits, experience, composition of the voting public and financial situation might influence the outcome. The question was raised if the web survey alone would be sufficient to reach a consensus for future meetings. Limiting a consensus meeting to only the websurvey would limit the time as well as the costs involved. Moreover, a higher percentage of surgeons participated in the web survey. In our opinion, however, the integration of an actual face-to-face meeting in the consensus methodology raises more awareness, provides an opportunity to discuss views, and encourages the transfer of knowledge eventually leading to the creation of a widely supported paper.

The literature review was ended in December 2014. No studies after that were integrated for the consensus meeting as this was decided in our methodology. Therefore, new studies might have been conducted on some topics. Future research should be focused on the laparoscopic appendectomy in pregnant patients, elucidating the value of MRI in specific patient groups, evaluating the outcomes of initial non-operative treatment for both uncomplicated and

complicated appendicitis, specific patient groups and the need for interval appendectomy. We therefore propose that these statements are updated on a regular basis.

Although some limitations can be identified in our methodology, we have integrated a new systematic method for a consensus meeting. In our opinion this is the way forward and we need to efflorescence this method. Reproducibility, involving members of the scientific community and applicability are key components of a consensus meeting. We believe that only after evaluation of the general opinion within the EAES such guidelines should be put into order.

In conclusion, the consensus meeting of the EAES resulted in several statements and recommendations regarding the diagnosis and management of appendicitis based upon available evidence and expert opinion and was supported by the European surgical community. It provides guidance to surgeons and surgical residents facing patients with acute appendicitis.

## References

1. Andersson R, Hugander A, Thulin A, et al (1994) Indications for operation in suspected appendicitis and incidence of perforation. *BMJ* 308:107-110.
2. Basoli A, Zarba Meli E, Salvio A, et al (1993) Trends in the incidence of acute appendicitis in Italy during the past 30 years. *Minerva Chir* 48:127-132.
3. Elangovan S, Knapp DP, Kallail KJ (1997) Incidence of acute appendicitis confirmed by histopathologic diagnosis. *Kans Med* 98:10-13.
4. Ilves I, Fagerström A, Herzig KH, et al (2014) Seasonal variations of acute appendicitis and nonspecific abdominal pain in Finland. *World J Gastroenterol* 20:4037-4042
5. Viniol A, Keunecke C, Biroga T et al (2014) Studies of the symptom abdominal pain--a systematic review and meta-analysis. *Fam Pract* 31:517-529.
6. Wei PL, Chen CS, Keller JJ, et al (2012) Monthly variation in acute appendicitis incidence: a 10-year nationwide population-based study. *J Surg Res* 178:670-676.
7. Oguntola AS, Adeoti ML, Oyemolade TA (2010) Appendicitis: Trends in incidence, age, sex, and seasonal variations in South-Western Nigeria. *Ann Afr Med* 9:213-217.
8. Petroianu A, Oliveira-Neto JE, Alberti LR (2004) Comparative incidence of acute appendicitis in a mixed population, related to skin color. *Arq Gastroenterol* 41:24-26.
9. Brink CF, Prinsloo H, van der Poel JS (1985) The seasonal incidence of acute appendicitis. *S Afr Med J* 68:156-158.
10. Blanco FC, Sandler AD, Nadler EP (2012) Increased incidence of perforated appendicitis in children with obesity. *Clin Pediatr (Phila)* 51:928-932.
11. Walker AR, Shipton E, Walker BF, et al (1989) Appendectomy incidence in black and white children aged 0 to 14 years with a discussion on the disease's causation. *Trop Gastroenterol* 10:96-101.
12. Fitz RH (1886) Perforating inflammation of the vermiform appendix. *Am J Med Sci* 92:321-346.
13. Bakker OJ, Go PM, Puylaert JB et al (2010) Guideline on diagnosis and treatment of acute appendicitis: imaging prior to appendectomy is recommended. *Ned Tijdschr Geneesk* 54:A303
14. Korndorffer JR Jr, Fellingner E, Reed W (2010) SAGES guideline for laparoscopic appendectomy. *Surg Endosc* 24:757-761.
15. Howick J (2009) Oxford Centre for Evidence-Based medicine – Levels of Evidence. Available at: <http://www.cebm.net/oxfords-centre-evidence-based-medicine-levels-evidence-march-2009>. March 2009. Accessed on the first of November 2014.
16. Guyatt GH, Oxman AD, Vist G, et al (2008) Rating quality of evidence and strength of recommendations GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 336:924–926.
17. Goldet G, Howick J (2013) Understanding GRADE: an introduction. *J Evid Based Med* 6:50-54.
18. Atkins D, Best D, Briss PA, et al (2004) Grading quality of evidence and strength of recommendations. *BMJ* 328:1490.
19. Scottish Intercollegiate Guidelines Network (SIGN) (1993). Sign methodology. Available at: <http://www.sign.ac.uk/methodology/index.html>. Accessed first of November 2014
20. Boukdedid R, Abdoul H, Loustau M, et al (2011) Using and Reporting the Delphi Method for Selecting Healthcare Quality Indicators: A Systematic Review. *PLoS ONE* 6, DOI: doi: 10.1371/journal.pone.0020476, June 9, 2011
21. Dalkey NC, Helmer O (1963) An experimental application of the Delphi method to the use of experts. *Management Science* 9:458-467.

22. Hasson F, Keeney S, McKenna H (2000) Research guidelines for the Delphi survey technique. *J Adv Nurs* 32:1008–1015.
23. Bundy DG, Byerley JS, Liles EA, et al (2007) Does this child have appendicitis? *JAMA* 298:438-451.
24. Andersson RE (2004) Meta-analysis of the clinical and laboratory diagnosis of appendicitis. *Br J Surg* 91:28-37.
25. Alvarado A (1986) A practical score for the early diagnosis of acute appendicitis. *Ann Emerg Med* 15:557-564.
26. Samuel M (2002) Pediatric appendicitis score. *J Pediatr Surg* 37:877-881.
27. Ohle R, O'Reilly F, O'Brien KK, et al (2011) The Alvarado score for predicting acute appendicitis: a systematic review. *BMC Med* DOI: 10.1186/1741-7015-9-139, December 28,2011
28. Kulik DM, Uleryk EM, Maguire JL (2013) Does this child have appendicitis? A systematic review of clinical prediction rules for children with acute abdominal pain. *J Clin Epidemiol* 66:95-104.
29. Ebell MH, Shinholser J (2014) What Are the Most Clinically Useful Cutoffs for the Alvarado and Pediatric Appendicitis Scores? A Systematic Review. *Ann Emerg Med* 64:365-372.
30. Carroll PJ, Gibson D, El-Faedy O, et al (2013) Surgeon-performed ultrasound at the bedside for the detection of appendicitis and gallstones: systematic review and meta-analysis. *Am J Surg* 205:102-108.
31. Douglas CD, Macpherson NE, Davidson PM, et al (2000) Randomised controlled trial of ultrasonography in diagnosis of acute appendicitis, incorporating the Alvarado score. *BMJ* 321:919-922.
32. Parker L, Nazarian LN, Gingold EL, et al (2014) Cost and radiation savings of partial substitution of ultrasound for CT in appendicitis evaluation: a national projection. *AJR Am J Roentgenol* 202:124-135.
33. Doria AS, Moineddin R, Kellenberger CJ, et al (2006) US or CT for Diagnosis of Appendicitis in Children and Adults? A Meta-Analysis. *Radiology* 241:83-94.
34. Horton MD, Counter SF, Florence MG, et al (2000) A prospective trial of computed tomography and ultrasonography for diagnosing appendicitis in the atypical patient. *Am J Surg* 179:379-381.
35. Kwok MY, Kim MK, Gorelick MH (2004) Evidence-based approach to the diagnosis of appendicitis in children. *Pediatr Emerg Care* 20:690-698.
36. van Randen A, Bipat S, Zwinderman AH, et al (2008) Acute appendicitis: meta-analysis of diagnostic performance of CT and graded compression US related to prevalence of disease. *Radiology* 249:97-106.
37. van Randen A, Laméris W, van Es HW, et al (2011) A comparison of the accuracy of ultrasound and computed tomography in common diagnoses causing acute abdominal pain. *Eur Radiol* 21:1535-1545.
38. Weston AR, Jackson TJ, Blamey S (2005) Diagnosis of appendicitis in adults by ultrasonography or computed tomography: a systematic review and meta-analysis. *Int J Technol Assess Health Care* 21:368-379.
39. Terasawa T, Blackmore CC, Bent S, et al (2004) Systematic review: computed tomography and ultrasonography to detect acute appendicitis in adults and adolescents. *Ann Intern Med* 141:537-546.
40. Hlibczuk V, Dattaro JA, Jin Z, et al (2010) Diagnostic accuracy of noncontrast computed tomography for appendicitis in adults: a systematic review. *Ann Emerg Med* 55:51-59.
41. Howell JM, Eddy OL, Lukens TW, et al (2010) Clinical policy: Critical issues in the evaluation and management of emergency department patients with suspected appendicitis. *Ann Emerg Med* 55:71-116.

42. Kaiser S, Frenckner B, Jorulf HK (2002) Suspected appendicitis in children: US and CT--a prospective randomized study. *Radiology* 223:633-638.
43. Neumayer L, Kennedy A (2003) Imaging in appendicitis: a review with special emphasis on the treatment of women. *Obstet Gynecol* 102:1404-1409.
44. Stephen AE, Segev DL, Ryan DP, et al (2003) The diagnosis of acute appendicitis in a pediatric population: to CT or not to CT. *J Pediatr Surg* 38:367-371.
45. Brenner D, Elliston C, Hall E, et al (2001) Estimated risks of radiation-induced fatal cancer from pediatric CT. *AJR Am J Roentgenol* 176:289-296.
46. Brenner DJ, Elliston CD, Hall EJ, et al (2001) Estimates of the cancer risks from pediatric CT radiation are not merely theoretical: comment on "point/counterpoint: in x-ray computed tomography, technique factors should be selected appropriate to patient size. against the proposition". *Med Phys* 28:2387-2388.
47. Kim K, Kim YH, Kim SY, et al (2012) Low-dose abdominal CT for evaluating suspected appendicitis. *N Engl J Med* 366:1596-1605.
48. Anderson BA, Salem L, Flum DR (2005) A systematic review of whether oral contrast is necessary for the computed tomography diagnosis of appendicitis in adults. *Am J Surg* 190:474-478.
49. Fonseca AL, Schuster KM, Kaplan LJ, et al (2014) The use of magnetic resonance imaging in the diagnosis of suspected appendicitis in pregnancy: shortened length of stay without increase in hospital charges. *JAMA Surg* 149:687-693.
50. Barger RL Jr, Nandalur KR (2010) Diagnostic performance of magnetic resonance imaging in the detection of appendicitis in adults: a meta-analysis. *Acad Radiol* 17:1211-1216.
51. Anderson SW, Rhea JT, Milch HN, et al (2010) Influence of body habitus and use of oral contrast on reader confidence in patients with suspected acute appendicitis using 64 MDCT. *Emerg Radiol* 17:445-453.
52. Sauerland S, Jaschinski T, Neugebauer EA (2010) Laparoscopic versus open surgery for suspected appendicitis. *Cochrane Database Syst Rev* 10:doi: 10.1002/14651858.CD001546, October 6, 2010.
53. Sohn M, Hoffmann M, Hochrein A, et al (2015) Laparoscopic Appendectomy Is Safe: Influence of Appendectomy Technique on Surgical-site Infections and Intra-abdominal Abscesses. *Surg Laparosc Endosc Percutan Tech* 25:e90-94.
54. Massoomi H, Mills S, Dolich MO, et al (2011) Comparison of outcomes of laparoscopic versus open appendectomy in adults: data from the nationwide inpatient sample (NIS), 2006-2008. *J Gastrointest Surg* 15:2226-2231.
55. Xiao Y, Shi G, Zhang J, et al (2014) Surgical site infection after laparoscopic and open appendectomy: a multicenter large consecutive cohort study. *Surg Endosc* 29:1384-1393.
56. Kocataş A, Gönenç M, Bozkurt MA, et al (2013) Comparison of open and laparoscopic appendectomy in uncomplicated appendicitis: a prospective randomized clinical trial. *Ulus Travma Acil Cerrahi Derg* 19:200-204.
57. Asarias JR, Schluskel AT, Cafasso DE, et al (2011) Incidence of postoperative intraabdominal abscesses in open versus laparoscopic appendectomies. *Surg Endosc* 25:2678-2683.
58. Othani H, Tamamori Y, Arimoto Y, et al (2012) Meta-analysis of the results of randomized controlled trials that compared laparoscopic and open surgery for acute appendicitis. *J Gastrointest Surg* 16:1929-1939.
59. Wilson DG, Bond AK, Ladwa N, et al (2013) Intra-abdominal collections following laparoscopic versus open appendectomy: an experience of 516 consecutive cases at a district general hospital. *Surg Endosc* 27:2351-2356.
60. Swank HA, Eshuis EJ, van Berge Henegouwen MI, et al (2011) Short- and long-term results of open versus laparoscopic appendectomy. *World J Surg* 35:1221-1226.

61. Li X, Zhang J, Sang L, et al (2010) Laparoscopic versus conventional appendectomy--a meta-analysis of randomized controlled trials. *BMC Gastroenterol*, DOI: 10.1186/1471-230X-10-129, November 3, 2010
62. Liu Z, Zhang P, Ma Y, et al (2010) Laparoscopy or not: A meta analysis of the surgical effects of laparoscopic versus open appendicectomy. *Surg Laparosc Endosc Percutan Tech* 20:362-370.
63. Wei B, Qi CL, Chen TF, et al (2011) Laparoscopic versus open appendectomy for acute appendicitis: a metaanalysis. *Surg Endosc* 25:1199-1208.
64. Wei HB, Huang JL, Zheng ZH, et al (2010) Laparoscopic versus open appendectomy: a prospective randomized comparison. *Surg Endosc* 24:266-269.
65. Markar SR, Penna M, Harris A (2014) Laparoscopic approach to appendectomy reduces the incidence of short- and long-term post-operative bowel obstruction: systematic review and pooled analysis. *J Gastrointest Surg* 18:1683-1692.
66. Kapischke M, Friedrich F, Hedderich J, et al (2011) Laparoscopic versus open appendectomy quality of life 7 years after surgery. *Langenbecks Arch Surg* 396:69-75.
67. Pelosi MA, Pelosi MA 3<sup>rd</sup> (1992) Laparoscopic appendectomy using a single umbilical puncture (minilaparoscopy). *J Reprod Med* 37:588-594.
68. Antoniou SA, Koch OO, Antoniou GA, et al (2014) Meta-analysis of randomized trials on single-incision laparoscopic versus conventional laparoscopic appendectomy. *Am J Surg* 207:613-622.
69. Frutos D, Abrisqueta J, Lujan J, et al (2013) Randomized prospective study to compare laparoscopic appendectomy versus umbilical single incision appendectomy. *Ann Surg* 257:413-418.
70. Gill RS, Shi X, Al-adra DP, et al (2012) Single-incision appendectomy is comparable to conventional laparoscopic appendectomy: A systematic review and pooled analysis. *Surg Laparosc Endosc Percutan Tech* 22:319-327.
71. Cai YL, Xiong XZ, Wu SJ, et al (2013) Single-incision laparoscopic appendectomy vs conventional laparoscopic appendectomy: systematic review and meta-analysis. *World J Gastroenterol* 19:5165-5173.
72. Clerveus M, Morandeira-Rivas A, Moreno-Sanz C, et al (2014) Systematic review and meta-analysis of randomized controlled trials comparing single incision versus conventional laparoscopic appendectomy. *World J Surg* 38:1937-1946.
73. Markar SR, Karthikesalingam A, Di Franco F, et al (2013) Systematic review and meta-analysis of single-incision versus conventional multiport appendectomy. *Br J Surg* 13:1709-1718.
74. Xu AM, Huang L, Li TJ (2015) Single-incision versus three-port laparoscopic appendectomy for acute appendicitis: systematic review and meta-analysis of randomized controlled trials. *Surg Endosc* 29:822-843.
75. Xue C, Lin B, Huang Z, et al (2015) Single-incision laparoscopic appendectomy versus conventional 3-port laparoscopic appendectomy for appendicitis: an updated meta-analysis of randomized controlled trials. *Surg Today* 45:1179-1186.
76. Zhou H, Jin K, Zhang J, et al (2014) Single incision versus conventional multiport laparoscopic appendectomy: a systematic review and meta-analysis of randomized controlled trials. *Dig Surg* 31:384-391.
77. Li P, Chen ZH, Li QG, et al (2013) Safety and efficacy of single-incision laparoscopic surgery for appendectomies: a meta-analysis. *World J Gastroenterol* 19:4072-4082.
78. Liang HH, Hung CS, Wang W, et al (2014) Single-incision versus conventional laparoscopic appendectomy in 688 patients: a retrospective comparative analysis. *Can J Surg* 57:E89-97.
79. Gao J, Li P, Li Q, et al (2013) Comparison between single-incision and conventional three-port laparoscopic appendectomy: a meta-analysis from eight RCTs. *Int J Colorectal Dis* 28:1319-1327.
80. Teoh AYW, Chiu PWY, Wong TCL, et al (2012) A double blinded randomized controlled trial of laparoendoscopic single site access versus conventional 3-port appendectomy. *Ann Surg* 256:909-914.

81. Lee WS, Choi ST, Lee JN, et al (2013) Single-Port laparoscopic appendectomy versus conventional laparoscopic appendectomy. A prospective randomized controlled study. *Ann Surg* 257:214-218.
82. Gorter RR, Heij HA, Eker HH, et al (2014) Laparoscopic appendectomy: State of the art. Tailored approach to the application of laparoscopic appendectomy? *Best Pract Res Clin Gastroenterol* 28:211-224.
83. Hua J, Gong J, Xu B, et al (2014) Single-incision versus conventional laparoscopic appendectomy: a meta-analysis of randomized controlled trials. *J Gastrointest Surg* 18:426-436.
84. Hansson J, Körner U, Khorram-Manesh A, et al (2009) Randomized clinical trial of antibiotic therapy versus appendectomy as primary treatment of acute appendicitis in unselected patients. *Br J Surg* 96:473-481.
85. Eriksson S, Granstrom L (1995) Randomized clinical trial of appendectomy versus antibiotic therapy for acute appendicitis. *Br J Surg* 82:166-169.
86. Styurd J, Eriksson S, Nilsson I, et al (2006) Appendectomy versus antibiotic treatment in acute appendicitis. A prospective multicentre randomized controlled trial. *World J Surg* 30:1033-1037.
87. Malik AA, Bari S (2009) Conservative management of acute appendicitis. *J Gastrointest Surg* 13:966-970.
88. Vons C, Barry C, Maitre S, et al (2011) Amoxicillin plus clavulanic acid versus appendicectomy for treatment of acute uncomplicated appendicitis. An open label, non-inferiority, randomised controlled trial. *Lancet* 377:1573-1579.
89. Wilms IMHA, de Hoog DENM, de Visser DC, et al (2011) Appendectomy versus antibiotic treatment for acute appendicitis (review) *Cochrane Database Syst Rev* DOI: 10.1002/14651858.CD008359. pub2, November 9, 2011
90. Svensson JF, Hall NJ, Eaton S, et al (2012) A review of conservative treatment of acute appendicitis. *Eur J Pediatr Surg* 22:185-194.
91. Mason RJ, Moazzez A, Sohn H, et al (2012) Meta-analysis of randomized trials comparing antibiotic therapy with appendectomy for acute uncomplicated (no abscess or phlegmon) appendicitis. *Surg Infect (Larchmt)* 13:74-84.
92. Kirby A, Hobson RP, Burke D, et al (2015) Appendicectomy for suspected uncomplicated appendicitis is associated with fewer complications than conservative antibiotic management: a meta-analysis of post-intervention complications. *J Infect* 70:105-110.
93. Varadhan KK, Neal KR, Lobo DN (2012) Safety and efficacy of antibiotics compared with appendicectomy for treatment of uncomplicated acute appendicitis: meta analysis of randomised controlled trials. *BMJ*, DOI: 10.1136/bmj.e2156, April 5, 2012.
94. Ansaloni L, Catena F, Coccolini F, et al (2011) Surgery versus conservative antibiotic treatment in acute appendicitis: a systematic review and meta-analysis of randomized controlled trials. *Dig Surg* 28:210-221.
95. Liu K, Fogg L (2011) Use of antibiotics alone for treatment of uncomplicated acute appendicitis: a systematic review and meta-analysis. *Surgery* 150:673-683.
96. Dimitriou I, Reckmann B, Nephuth O, et al (2013) Single institution's experience in laparoscopic appendectomy as a suitable therapy for complicated appendicitis. *Langenbecks Arch Surg* 398:147-152.
97. Thomson JE, Kruger D, Jann-Kruger C, et al (2015) Laparoscopic versus open surgery for complicated appendicitis: a randomized controlled trial to prove safety. *Surg Endosc* 29:2027-2032.
98. Tuggle KR, Ortega G, Bolorunduro OB, et al (2010) Laparoscopic versus open appendectomy in complicated appendicitis: a review of the NSQIP database. *J Surg Res* 163:225-228.
99. Andersson RE, Petzold MG (2007) Non-surgical treatment of appendiceal abscess or phlegmon. A systematic review and meta-analysis. *Ann Surg* 246:741-748.

100. Simillis C, Symeonides P, Shorthouse AJ, et al (2010) A meta-analysis comparing conservative treatment versus acute appendectomy for complicated appendicitis (abscess or phlegmon). *Surgery* 147:818-29.
101. Kim JK, Ryou S, Oh HK, et al (2010) Management of appendicitis presenting with abscess or mass. *J Korean Soc Coloproctol* 26:413-419.
102. Deelder JD, Richir MC, Schoorl T, et al (2014) How to treat an appendiceal inflammatory mass: operatively or nonoperatively? *J Gastrointest Surg* 18:641-645.
103. Hall NJ, Jones CE, Eaton S, et al (2011) Is interval appendectomy justified after successful nonoperative treatment of an appendix mass in children? A systematic review. *J Pediatr Surg* 46:767-771.
104. Clarke T, Katkhouda N, Mason RJ, et al (2011) Laparoscopic versus open appendectomy for the obese patient: a subset analysis from a prospective, randomized, double-blind study. *Surg Endosc* 25:1276-1280.
105. Markar SR, Venkat-Raman V, Ho A, et al (2011) Laparoscopic versus open appendectomy in obese patients. *Int J Surg* 9:451-455.
106. Woodham BL, Cox MR, Eslick GD (2012) Evidence to support the use of laparoscopic over open appendectomy for obese individuals: a meta-analysis. *Surg Endosc* 26:2566-2570.
107. Dasari BV, Baker J, Markar S, et al (2015) Laparoscopic appendectomy in obese is associated with improvements in clinical outcome: systematic review. *Int J Surg* 13:250-256.
108. Walsh CA, Tang T, Walsh SR (2008) Laparoscopic versus open appendectomy in pregnancy: A systematic review. *Int J Surg* 6:339-344.
109. Wilasrusmee C, Sukrat B, McEvoy M, et al (2012) Systematic review and meta-analysis of safety of laparoscopic versus open appendectomy for suspected appendicitis in pregnancy. *Br J Surg* 99:1470-1478.
110. McGory ML, Zingmond DS, Tillou A, et al (2007) Negative appendectomy in pregnant women is associated with substantial risk of fetal loss. *J Am Coll Surg* 205:534-540.
111. Walker HG, Al Samaraee A, Mills SJ, et al (2014) Laparoscopic appendectomy in pregnancy: a systematic review of the published evidence. *Int J Surg* 12:1235-1241.
112. Chung JC, Cho GS, Shin EJ, et al (2013) Clinical outcomes compared between laparoscopic and open appendectomy in pregnant women. *Can J Surg* 56:341-346.
113. Markar SR, Blackburn S, Cobb R, et al (2012) Laparoscopic versus open appendectomy for complicated and uncomplicated appendicitis in children. *J GastrointestSurg* 16:1993-2004.
114. Zwintscher NP, Johnson EK, Martin MJ, et al (2013) Laparoscopy utilization and outcomes for appendicitis in small children. *J Pediatr Surg* 48:1941-1945.
115. Armstrong J, Merritt N, Jones S, et al (2014) Non-operative management of early, acute appendicitis in children: is it safe and effective? *J Pediatr Surg* 49:782-785.
116. Svensson JF, Patkova B, Almström M, et al (2015) Nonoperative treatment with antibiotics versus surgery for acute nonperforated appendicitis in children: a pilot randomized controlled trial. *Ann Surg* 261:67-71.
117. Yeh CC, Wu SC, Liao CC, et al (2011) Laparoscopic appendectomy for acute appendicitis is more favorable for patients with comorbidities, the elderly, and those with complicated appendicitis: a nationwide population-based study. *Surg Endosc* 25:2932-2942.
118. Masoomi H, Mills S, Dolich MO, et al (2012) Does laparoscopic appendectomy impart an advantage over open appendectomy in elderly patients? *World J Surg* 36:1534-1539.
119. Southgate E, Vousden N, Karthikesalingam A, et al (2012) Laparoscopic vs open appendectomy in older patients. *Arch Surg* 147:557-562.
120. Wang YC, Yang HR, Chung PK, et al (2006) Laparoscopic appendectomy in the elderly. *Surg Endosc* 20:887-889.

121. Kim MJ, Fleming FJ, Gunzler DD, et al (2011) Laparoscopic appendectomy is safe and efficacious for the elderly: an analysis using the National Surgical Quality Improvement Project database. *Surg Endosc* 25:1802-1807.
122. McCartan DP, Fleming FJ, Grace PA (2010) The management of right iliac fossa pain - Is timing everything? *Surgeon* 8:211-217.
123. Fair BA, Kubasiak JC, Janssen I, et al (2015) The impact of operative timing on outcomes of appendicitis: a National Surgical Quality Improvement Project analysis. *Am J Surg* 209:498-502.
124. Salminen P, Paajanen H, Rautio T, et al (2015) Antibiotic Therapy vs Appendectomy for Treatment of Uncomplicated Acute Appendicitis: The APPAC Randomized Clinical Trial. *JAMA* 313:2340-2348.
125. Cappendijk VC, Hazebroek FW (2000) The impact of diagnostic delay on the course of acute appendicitis. *Arch Dis Child* 83:64-66.
126. Papandria D, Goldstein SD, Rhee D, et al (2013) Risk of perforation increases with delay in recognition and surgery for acute appendicitis. *J Surg Res* 184:723-729.
127. United Kingdom National Surgical Research Collaborative, Bhangu A (2014) Safety of Short, In-Hospital Delays Before Surgery for Acute Appendicitis: Multicentre Cohort Study, Systematic Review, and Meta-Analysis. *Ann Surg* 259:894-903.
128. Shin CS, Roh YN, Kim JL (2014) Delayed appendectomy versus early appendectomy in the treatment of acute appendicitis: a retrospective study. *World J Emerg Surg* DOI: 10.1186/1749-7922-9-8, January 21, 2014.
129. Ingraham AM, Cohen ME, Bilimoria KY, et al (2010) Effect of delay of operation on outcomes in adults with acute appendicitis. *Arch Surg* 145: 886-892.
130. Andersen BR, Kallehave FL, Andersen HK (2005) Antibiotics versus placebo for prevention of postoperative infection after appendectomy. *Cochrane Database Syst Rev* DOI: 10.1002/14651858.CD001439.pub2, July 20, 2005.
131. Daskalakis K, Juhlin C, Pählman L (2014) The use of pre- or postoperative antibiotics in surgery for appendicitis: A systematic review. *Scand J Surg* 103:14-20.
132. Kasatpibal N, Nørgaard M, Sørensen HT, et al (2006) Risk of surgical site infection and efficacy of antibiotic prophylaxis: a cohort study of appendectomy patients in Thailand. *BMC Infect Dis*, DOI: 10.1186/1471-2334-6-111, July 12, 2006.
133. Angioli R, Terranova C, De Cicco Nardone C, et al (2013) A comparison of three different entry techniques in gynecological laparoscopic surgery: a randomized prospective trial. *Eur J Obstet Gynecol Reprod Biol* 171:339-342.
134. Agresta F, De Simone P, Ciardo LF, et al (2004) Direct trocar insertion vs Veress needle in nonobese patients undergoing laparoscopic procedures: a randomized prospective single-center study. *Surg Endosc* 18:1778-1781.
135. Bemelman WA, Dunker MS, Busch OR, et al (2000) Efficacy of establishment of pneumoperitoneum with the Veress needle, Hasson trocar, and modified blunt trocar (TrocDoc): a randomized study. *J Laparoendosc Adv Surg Tech A* 10:325-330.
136. van Rossem CC, Bolmers MD, Schreinemacher MH, et al (2016) Prospective nationwide outcome audit of surgery for suspected acute appendicitis. *Br J Surg* 103:144-151.
137. Hamminga JT, Hofker HS, Broens PM, et al (2013) Evaluation of the appendix during diagnostic laparoscopy, the laparoscopic appendicitis score: a pilot study. *Surg Endosc* 27:1594-1600.
138. Strong S, Blencowe N, Bhangu A, et al (2015) How good are surgeons at identifying appendicitis? Results from a multi-centre cohort study. *Int J Surg* 15:107-112.
139. Champault G, Rizk N, Ziol M, et al (1996) Can we recognize the pathological character of the appendix during laparoscopy? Prospective study: 81 cases. *J Chir (Paris)* 133:320-323.
140. Grunewald B, Keating J (1993) Should the 'normal' appendix be removed at operation for appendicitis? *J R Coll Surg Edinb* 38:158-160.

141. Sajid MS, Rimple J, Cheek E, et al (2009) Use of endo-GIA versus endo-loop for securing the appendicular stump in laparoscopic appendectomy: a systematic review. *Surg Laparosc Endosc Percutan Tech* 19:11-15.
142. Ates M, Dirican A, Ince V, et al (2012) Comparison of Intracorporeal Knot-tying Suture (Polyglactin) and Titanium Endoclips in Laparoscopic Appendiceal Stump Closure. *Surg Laparosc Endosc Percutan Tech* 22:226-231.
143. Kazemier G, In't Hof KH, Saad S, et al (2006) Securing the appendiceal stump in laparoscopic appendectomy: Evidence for routine stapling? *Surg Endosc* 20:1473-1476.
144. St Peter SD, Adibe OO, Iqbal CW, et al (2012) Irrigation Versus Suction Alone During Laparoscopic Appendectomy for Perforated Appendicitis. *Ann Surg* 256:581-585.
145. Akkoyun I, Tuna AT (2012) Advantages of abandoning abdominal cavity irrigation and drainage in operations performed on children with perforated appendicitis. *J Pediatr Surg* 47:1886-1890.
146. Hartwich JE, Carter RF, Wolfe L, et al (2013) The effects of irrigation on outcomes in cases of perforated appendicitis in children. *J Surg Res* 180:222-225.
147. Allemann P, Probst H, Demartines N, et al (2011) Prevention of infectious complications after laparoscopic appendectomy for complicated acute appendicitis-the role of routine abdominal drainage. *Langenbecks Arch Surg* 396:63-68.
148. St Peter SD, Aguayo P, Fraser JD, et al (2010) Initial laparoscopic appendectomy versus initial nonoperative management and interval appendectomy for perforated appendicitis with abscess: a prospective, randomized trial. *J Pediatr Surg* 45:236-240.
149. Olsen J, Skovdal J, Qvist N, et al (2014) Treatment of appendiceal mass - A qualitative systematic review. *Dan Med J* 61:A4881.
150. Misdraji J, Yantiss RK, Graeme-Cook FM, et al (2003) Appendiceal mucinous neoplasms: a clinicopathologic analysis of 107 cases. *Am J Surg Pathol* 27:1089-1103.
151. McDonald JR, O'Dwyer ST, Rout S, et al (2012) Classification of and cytoreductive surgery for low-grade appendiceal mucinous neoplasms. *Br J Surg* 99:987-992.
152. Arnason T, Kamionek M, Yang M, et al (2015) Significance of proximal margin involvement in low-grade appendiceal mucinous neoplasms. *Arch Pathol Lab Med* 139:518-521.
153. Henderson L, Fehily C, Folaranmi S, et al (2014) Management and outcome of neuroendocrine tumours of the appendix-a two centre UK experience. *J Pediatr Surg* 49:1513-1517.
154. Kim SS, Kays DW, Larson SD, et al (2014) Appendiceal carcinoids in children--management and outcomes. *J Surg Res* 192:250-253.
155. Coakley B, Sussman E, Wolfson T, et al (2011) Postoperative antibiotics correlate with worse outcomes after appendectomy for nonperforated appendicitis. *J Am Coll Surg* 213:778-783.
156. van Rossem CC, Schreinemacher M, Treskes K, et al (2014) Duration of antibiotic treatment after appendectomy for acute complicated appendicitis. *Br J Surg* 101: 715-719.
157. Pinto DJ, Sanderson PJ (1980) Rational use of antibiotic therapy after appendectomy. *Br J Surg* 280:275-277.
158. Ambrose NS, Donovan IA, Wise R, et al (1983) Metronidazole and ticarcillin in the prevention of sepsis after appendectomy. *Am J Surg* 146:346-348.
159. Campbell WB (1980) Prophylaxis of infection after appendectomy: a survey of current surgical practice. *BMJ* 281:1597-1600.
160. Buckels JA, Brookstein R, Bonser R, et al (1985) A comparison of the prophylactic value of cefotetan and metronidazole appendectomy. *World J Surg* 9:814-818.
161. Nadler E, Reblock KK, Ford HR, et al (2003) Monotherapy versus multi-drug therapy for the treatment of perforated appendicitis in children. *Surg Infect (Larchmt)* 4:327-333.
162. Yu TC, Hamill JKM, Evans SM, Price NR, et al (2013) Duration of Postoperative Intravenous Antibiotics in Childhood Complicated Appendicitis: A Propensity Score-Matched Comparison Study. *Eur J Pediatr Surg* 24:341-349.

163. Lau WY, Fan ST, Yiu TF, et al (1983) Prophylaxis of post-appendectomy sepsis by metronidazole and ampicillin: a randomized, prospective and double-blind trial. *Br J Surg* 70:155-157.
164. Lau DHW, Yau KKK, Chung CC, et al (2005) Comparison of needlescopic appendectomy versus conventional laparoscopic appendectomy: a randomized controlled trial. *Surg Laparosc Endosc Percutan Tech* 15:75-79.
165. Andersson RE (2001) Small bowel obstruction after appendectomy. *Br J Surg* 88:1387-1391.
166. Emil S, Elkady S, Shbat L, et al (2014) Determinants of postoperative abscess occurrence and percutaneous drainage in children with perforated appendicitis. *Pediatr Surg Int* 30:1265-1271.
167. Fleming FJ, Kim MJ, Messing S, et al (2010) Balancing the risk of postoperative surgical infections: a multivariate analysis of factors associated with laparoscopic appendectomy from the NSQIP database. *Ann Surg* 252:895-900.
168. Bahar MM, Jangjoo A, Amouzesi A, et al (2010) Wound infection incidence in patients with simple and gangrenous or perforated appendicitis. *Arch Iran Med* 13:13-16.
169. Guller U, Hervey S, Purves H, et al (2004) Laparoscopic versus open appendectomy: outcomes comparison based on a large administrative database. *Ann Surg* 239:43-52.
170. Isaksson K, Montgomery A, Moberg AC, et al (2014) Long-term follow-up for adhesive small bowel obstruction after open versus laparoscopic surgery for suspected appendicitis. *Ann Surg* 259:1173-1177.
171. Boomer LA, Cooper JN, Deans KJ, et al (2014) Does delay in appendectomy affect surgical site infection in children with appendicitis? *J Pediatr Surg* 49:1026-1029.
172. Graat LJ, Bosma E, Roukema JA, et al (2012) Appendectomy by Residents Is Safe and Not Associated With a Higher Incidence of Complications. *Ann Surg* 255:715-719.
173. Gandaglia G, Ghani KR, Sood A, et al (2014) Effect of minimally invasive surgery on the risk for surgical site infections: results from the National Surgical Quality Improvement Program (NSQIP) Database. *JAMA Surg* 149:1039-1044.
174. Advani V, Ahad S, Gonczy C, et al (2012) Does resident involvement effect surgical times and complication rates during laparoscopic appendectomy for uncomplicated appendicitis? An analysis of 16,849 cases from the ACS-NSQIP. *Am J Surg* 203:347-351.
175. Galli R, Banz V, Fenner H, et al (2013) Laparoscopic approach in perforated appendicitis: increased incidence of surgical site infection? *Surg Endosc* 27:2928-2933.
176. Kanona H, Al Samaraee A, Nice C, et al (2012) Stump appendicitis: a review. *Int J Surg* 10:425-428
177. Clark JJ, Johnson SM (2011) Laparoscopic drainage of intraabdominal abscess after appendectomy: an alternative to laparotomy in cases not amenable to percutaneous drainage. *J Pediatr Surg* 46:1385-1389.
178. Ben Dhaou M, Ghorbel S, Chouikh T, et al (2010) Conservative management of post-appendectomy intra-abdominal abscesses. *Ital J Pediatr* DOI:10.1186/1824-7288-36-68, October 14, 2010.
179. Forgues D, Habbig S, Dillo AF, et al (2007) Post-appendectomy intra-abdominal abscesses; Can they successfully be managed with the sole use of antibiotic therapy? *Eur J Pediatr Surg*;17:104-109
180. Contreras-Dominguez V, Carbonell-Bellilio C (2008) Profylactic antiemetic therapy for acute abdominal surgery. A comparative study of Droperidol, Metocloperamide, Tropisetron, granisetron and Dexamethasone. *Rev Bras Anesthesiol* 58:35-44.
181. Morton NS, K O'Brien K (1999) Analgesic efficacy of paracetamol and diclofenac in children receiving PCA morphine. *Br J Anaesth* 82:715-717.
182. Jalil RMA, Yahya N, Sulaimn O, et al (2013) Comparing the effectiveness of ropivacaine 0.5% versus ropivacaine 0.2% for transabdominis plane block in providing postoperative analgesia after appendectomy. *Acta Anaesthesiol Taiwan* 52:49-53.

183. Ahn SR, Kang DB, Lee C, et al (2013) Postoperative pain relief using wound infiltration with 0.5% bupivacaine in single-incision laparoscopic surgery for an appendectomy. *Ann Coloproctol* 29:238-242.
184. Cho S, Kim YJ, Kim DY, et al (2013) Postoperative analgesic effects of ultrasound-guided transversus abdominis plane block for open appendectomy. *J Korean Surg Soc* 85:128-133.
185. Taal BG (2005) Diagnosis and treatment in intestinal carcinoid tumours. *Curr Gastroenterol Rep* 7:1-3.
186. Sieren LM, Collins JN, Weireter LJ, et al (2010) The incidence of benign and malignant neoplasia presenting as acute appendicitis. *Am Surg* 76:808-811.
187. Benedix F, Reimer A, Gastinger I, et al (2010) Primary appendiceal carcinoma—epidemiology, surgery and survival: results of a German multi-center study. *Eur J Surg Oncol* 36:763-71.
188. Charfi S, Sellami A, Affes A, et al (2014) Histopathological findings in appendectomy specimens: A study of 24,697 cases. *Int J Colorectal Dis* 29:1009-1012.
189. Anwar K, Desai M, Al-Bloushi N, et al (2014) Prevalence and clinicopathological characteristics of appendiceal carcinoids in Sharjah (United Arab Emirates). *World J Gastrointest Oncol* 6:253-256
190. In 't Hof K, Wal HC, Kazemier G, et al (2008) Carcinoid tumour of the appendix: An analysis of 1,485 consecutive emergency appendectomies. *J Gastrointest Surg* 12:1436-1438.
191. Chandrasegaram MD, Rothwell LA, An EI, et al (2012) Pathologies of the appendix: A 10-year review of 4670 appendectomy specimens. *ANZ J Surg* 82:844-847.
192. Ramezani MA, Dehghani MR (2007) Relationship between *Enterobius vermicularis* and the incidence of acute appendicitis. *Southeast Asian J Trop Med Public health* 38:20-23.
193. Chong VH, Telisinghe PU, Yapp KS, et al (2011) Tuberculous appendix: A review of clinical presentations and outcomes. *Singapore Med J* 52:90-93.
194. Khan SA, Khokhar HA, Nasr AR, et al (2013) Incidence of right-sided colonic tumours (non-appendiceal) in patient's 40 years of age presenting with features of acute appendicitis. *Int J Surg* 11:301-304.
195. González-Moreno S, Sugarbaker PH (2004) Right hemicolectomy does not confer a survival advantage in patients with mucinous carcinoma of the appendix and peritoneal seeding. *Br J Surg* 91:304-311.
196. Amini A, Masoumi-Moghaddam S, Ehteda A, et al (2014) Secreted mucins in pseudomyxoma peritonei: pathophysiological significance and potential therapeutic prospects. *Orphanet J Rare Dis*, DOI: 10.1186/1750-1172-9-71, May 5, 2014.
197. Shapiro R, Eldar S, Sadot E, et al (2011) Appendiceal carcinoid at a large tertiary center: Pathologic findings and long-term follow-up evaluation. *Am J Surg* 201:805-808.
198. Marudanayagam R, Williams GT, Rees BI (2006) Review of the pathological results of 2660 appendectomy specimens. *J Gastroenterol* 41:745-749.
199. Connor SJ, Hanna GB, Frizelle FA (1997) Appendiceal Tumors: retrospective clinicopathologic analysis of appendiceal tumours from 7,970 Appendectomies. *Dis Colon Rectum* 41:75-80.
200. Alemayehu H, Snyder CL, St Peter SD, et al (2014) Incidence and outcomes of unexpected pathology findings after appendectomy. *J Pediatr Surg* 49:1390-1393.
201. Lobo-machín I, Delgado-Plasencia L, Hernández-González I, et al (2014) Appendiceal diverticulitis and acute appendicitis: differences and similarities. *Rev Esp Enferm Dig* 106:452-458.
202. Lohsiriwat V, Vongjirad A, Lohsiriwat D (2009) Incidence of synchronous appendiceal neoplasm in patients with colorectal cancer and its clinical significance. *World J Surg Oncol*, DOI: 10.1186/1477-7819-7-51, June 2, 2009.
203. Vettoretto N, Gobbi S, Corradi A, et al (2011) Consensus conference on laparoscopic appendectomy: development of guidelines. *Colorectal Dis* 13:748-754.

*Chapter 4*

204. Nair R, Aggarwal R, Khanna D (2011) Methods of formal consensus in classification/diagnostic criteria and guideline development. *Semin Arthritis Rheum* 41:95-105.
205. Rycroft-Malone J (2001) Formal consensus: the development of a national clinical guideline. *Qual Health Care* 10:238-44.
206. Loblaw DA, Prestrud AA, Somerfield MR, et al (2012) American Society of Clinical Oncology Clinical Practice Guidelines: formal systematic review-based consensus methodology. *J Clin Oncol* 30:3136-3140.