

Contents

- 1 Introduction 3**
 - 1.1 General introduction 3
 - 1.1.1 Relevance of the topic 3
 - 1.2 Background 4
 - 1.3 Initial Hypothesis 6
 - 1.3.1 Boreal wetlands 6
 - 1.3.2 Tropical wetlands 7
 - 1.3.3 Clathrates 7
 - 1.3.4 Geological sources 8
 - 1.4 Current state of research 8
 - 1.4.1 Paleo-environments 8
 - 1.4.2 Modelling CH₄ 9
 - 1.5 Aims and objectives 10
 - 1.6 Thesis approach and outline 11

- 2 Uncertainties: effect of hydrological model and structure 15**
 - 2.1 Abstract 15
 - 2.2 Introduction 16
 - 2.3 Models 17
 - 2.3.1 Previous modelling experiments 17
 - 2.3.2 Modelling experiments in this study 18
 - 2.4 Modelled climate changes 20
 - 2.4.1 Paleogeography: climate and environment 20
 - 2.4.2 Paleogeography: wetland distribution 22
 - 2.4.3 Present climate and environment and model validation 22
 - 2.5 Results 25
 - 2.5.1 Comparison of modelled values with data for MODERN climate 25
 - 2.5.2 Sensitivity to CH₄ production parameters in PEATLAND 25
 - 2.5.3 Sensitivity to hydrology and CH₄ emission model structure 28
 - 2.6 Discussion 30
 - 2.6.1 Model structure 30
 - 2.7 Conclusions 33

3	Uncertainties modelling CH₄ emissions from northern wetlands in glacial climates: the role of vegetation parameters	35
3.1	Abstract	35
3.2	Introduction	36
3.3	Effects of plants on CH ₄ fluxes	39
3.4	Modelling experiments	41
3.4.1	The PEATLAND-VU model	41
3.4.2	Climate model input to PEATLAND-VU	41
3.4.3	Model runs	42
3.5	Results	43
3.6	Discussion	46
3.7	Conclusions	49
4	Methane emission from permafrost thaw lakes limited by lake drainage	55
4.1	Abstract	55
4.2	Introduction	56
4.3	The model	56
4.3.1	Approach	56
4.3.2	Drivers and settings	56
4.4	Test and validation	57
4.4.1	The TERRAIN experiment	57
4.4.2	The ICEGROWTH experiment	57
4.4.3	The TIMING experiment	57
4.4.4	The FUTURE experiment	58
4.5	Methods	58
4.6	Results and discussion	60
5	Permafrost ecosystem: wetlands characteristics and their influence on CH₄ emissions in a drained thaw lake basin, northeastern Siberia	67
5.1	Abstract	67
5.2	Introduction	68
5.3	General settings of the site	70
5.4	Analytical techniques and methods	73
5.4.1	Soil analyses	73
5.4.2	CH ₄ oxidation rate analysis	74
5.4.3	DNA analysis	76
5.4.4	CH ₄ flux measurements	76
5.5	Results	77
5.5.1	Soils	77
5.5.2	CH ₄ oxidation rates	83
5.5.3	DNA	85
5.5.4	CH ₄ flux measurements	86
5.6	Discussion	87
5.7	Conclusions	90

6	Paleo-thaw lakes and their role as CH₄ sources	93
6.1	Abstract	93
6.2	Introduction	93
6.3	Study area	96
6.4	Methods	98
6.4.1	Cores and section logs	98
6.4.2	Grain size distribution analysis	99
6.4.3	Organic Carbon analysis	99
6.4.4	DNA analysis	99
6.4.5	Radiocarbon dating	99
6.5	Results	100
6.5.1	Cores and section logs	100
6.5.2	Grain size distribution analysis	104
6.5.3	Organic Carbon analysis	104
6.5.4	DNA analysis	108
6.5.5	Radiocarbon dating	109
6.5.6	Stratigraphy	109
6.5.7	Sedimentation rates	110
6.5.8	Weichselian environmental change recorded in the Hengelo area	110
6.5.9	Weichselian environmental change recorded in the Ootmarsum area	112
6.5.10	Weichselian environmental change recorded in the Orvelte exposure	113
6.6	Discussion	113
6.7	Conclusions	115
7	Conclusions	117
7.1	The initial questions	117
7.2	Main findings of this thesis	118
7.3	Current state of modelling northern wetland CH ₄ fluxes	119
7.4	Summary	119
7.4.1	Chapter 2	120
7.4.2	Chapter 3	121
7.4.3	Chapter 4	122
7.4.4	Chapter 5	123
7.4.5	Chapter 6	124
7.5	A critical look	125
7.6	Going forward	127
8	Nederlandse samenvatting	129
8.1	De onderzoeksvragen en hypothesen	129
8.2	Belangrijkste resultaten van dit onderzoek	130
8.3	Huidige status van modellering methaanfluxen	131
8.4	Samenvatting	132
8.4.1	Hoofdstuk 2	132
8.4.2	Hoofdstuk 3	134
8.4.3	Hoofdstuk 4	135
8.4.4	Hoofdstuk 5	136
8.4.5	Hoofdstuk 6	137

8.5 Een kritische blik 138
8.6 Een vooruitblik 139

Appendix **173**